

REVIEW ARTICLE

Parasitic helminths: a pharmacopeia of anti-inflammatory molecules

M. J. G. JOHNSTON^{1,2}, J. A. MACDONALD^{3,4} and D. M. MCKAY^{1,2*}¹Gastrointestinal Research Group, ²Department of Physiology and Biophysics, ³Smooth Muscle Research Group and ⁴Department of Biochemistry and Molecular Biology, University of Calgary, 3330 Hospital Drive NW, Calgary, Alberta, Canada, T2N 1N4

(Received 15 July 2008; revised 25 August and 22 September 2008; accepted 1 October 2008; first published online 15 December 2008)

SUMMARY

Infection with parasitic helminths takes a heavy toll on the health and well-being of humans and their domestic livestock, concomitantly resulting in major economic losses. Analyses have consistently revealed bioactive molecules in extracts of helminths or in their excretory/secretory products that modulate the immune response of the host. It is our view that parasitic helminths are an untapped source of immunomodulatory substances that, in pure form, could become new drugs (or models for drug design) to treat disease. Here, we illustrate the range of immunomodulatory molecules in selected parasitic trematodes, cestodes and nematodes, their impact on the immune cells in the host and how the host may recognize these molecules. There are many examples of the partial characterization of helminth-derived immunomodulatory molecules, but these have not yet translated into new drugs, reflecting the difficulty of isolating and fully characterizing proteins, glycoproteins and lipid-based molecules from small amounts of parasite material. However, this should not deter the investigator, since analytical techniques are now being used to accrue considerable structural information on parasite-derived molecules, even when only minute quantities of tissue are available. With the introduction of methodologies to purify and structurally-characterize molecules from small amounts of tissue and the application of high throughput immunological assays, one would predict that an assessment of parasitic helminths will yield a variety of novel drug candidates in the coming years.

Key words: helminths, immunomodulators, anti-inflammatory drugs, drug design.

INTRODUCTION

Infections with parasitic helminths continue to cause considerable morbidity and mortality in humans and their domestic livestock, translating into substantial socioeconomic losses. For instance, morbidity associated with *Schistosoma mansoni* or *Echinococcus granulosus* infections is debilitating and sometimes fatal (Carod-Artal, 2008; King and Dangerfield-Cha, 2008), and infections with hookworms retards the growth of children and can impair their cognitive functions (Oberhelman *et al.* 1998). The magnitude of the problems is illustrated by estimates from the the World Health Organization (WHO, 1999) that, even in the midst of successful anthelmintic programs, 30% of the world's population harbours at least 1 species of helminth parasite; a staggering 2 billion people are affected! However, numerous studies have shown that modulation of the immune

response of the host by parasitic helminths can have a concomitant health benefit (reviewed by McKay, 2006). In addition to infections with live organisms, helminth-derived molecules are potent immunomodulators and can serve as templates for the design of novel anti-inflammatory drugs, and may also be candidates for vaccine development (Harnett and Harnett, 2008; Perrigoue *et al.* 2008). The former aspect of the host-parasite interaction is the topic of this review.

HELMINTH THERAPY: THE GOOD, THE BAD AND THE WRIGGLY

Parasitic helminths have been humans' long-time companion and have evolved as 'master regulators' of host immune responses (Maizels *et al.* 2004). Consequently, knowledge of how the parasite modifies host immunity can be used to develop therapies for other diseases. In considering 'helminth therapy', one can either use viable infections or immunomodulatory molecules from the parasite; both strategies have been used to treat diseases in animal models. Infection with a variety of species of

* Corresponding author: Health Sciences Centre, HSC 1877, University of Calgary, 3330 Hospital Drive NW, Calgary, Alberta, Canada T2N 1N4. Tel: +1 403 220 7362. Fax: +1 403 283 3028. E-mail: dmckay@ucalgary.ca

parasitic helminths has been shown to reduce the severity of airway, cerebral and intestinal inflammation in murine models of human disease (McKay, 2006), and a number of investigators are pursuing the mechanism(s) underlying this health benefit (Weinstock *et al.* 2005). Moreover, Weinstock and colleagues have shown that the ingestion of viable eggs of *Trichuris suis* by patients with inflammatory bowel disease (Crohn's disease and ulcerative colitis) led to a reduction in disease activity, and adverse side-effects were not apparent (Summers *et al.* 2003, 2005). However, there can be a number of concerns regarding the use of an infection as a therapy. First, a parasite that has limited impact on the host must be used, which prohibits the use of particular species, such as *S. mansoni*, despite its potent stimulation of host immunity. Second, there are patient groups for which this approach would not be advocated, typically children, the elderly, immunocompromised individuals and pregnant women (Farah *et al.* 2007). Third, there are examples of helminth infections worsening disease evoked by bacteria or protozoa (Chen *et al.* 2005; Marshall *et al.* 1999) and, in at least one case, chemically-induced colitis (Hunter *et al.* 2007). Finally, when introducing a species to a novel environment, there is the potential for maladaptive responses and unpredicted pathological effects/changes in remote tissues. Moreover, the possibility exists for parasite establishment in, rather than expulsion from the host (Kradin *et al.* 2006). Despite these issues, the data on 'helminth therapy' are encouraging, and animal models can be used to identify novel anti-inflammatory mechanisms for translation into medical treatments, negating the need for actual infection.

The use of helminth-derived molecules as drugs overcomes the concerns with the use of live parasites. However, the issues here are often technical and revolve around a single basic question: how can pure molecules be isolated from miniscule amounts of parasite material to allow testing in biological systems and to serve as blueprints for drug development?

IMMUNE RESPONSES TO HELMINTH PARASITES: THE BASICS

A number of authoritative reviews of the mammalian immune system and the responses to infection with helminths are available (Hayes *et al.* 2004; Maizels *et al.* 2004; Antony *et al.* 2007; Khan, 2008; Perrigoue *et al.* 2008). As metazoans, helminths cannot be ingested by macrophages and other phagocytes, so that a variety of effector mechanisms have evolved to combat helminths. These involve the mobilization and activation of T and B lymphocytes, mast cells, eosinophils, natural killer (NK) cells, and, if the infection is at a mucosal surface, mucus-producing goblet cells. Briefly, while there is likely to be an

innate immune response to helminth-derived products by macrophages and natural killer cells, the current view is that the predominant anti-worm response is the remit of an adaptive immune response. Thus, following antigen processing and presentation by professional antigen presenting cells (e.g., dendritic cells (DC Φ s)), humans and mice display a stereotypic T cell response which is characterized by the development of CD4⁺ T helper (TH)-2 cells that produce interleukin (IL)-4, -5, -13 and others. These chemical signals mobilize other cells; for example, they direct isotype switching, so that high affinity immunoglobulin (Ig) G, E and A can be produced. These antibodies then bind to receptors on effector cells, allowing for their activation by worm antigen and the release of molecules (e.g., mast-derived histamine and trypases, eosinophil-derived cationic protein) that either disrupt the surface of the helminth and/or alter the environment, making it less hospitable to the helminth. In addition, IgG can bind directly to the surface of the helminth (opsonization), interfering with worm behaviour and targeting the worm for attack by complement and polymorphonuclear cells. Thus, the host response to infection with parasitic helminths is multi-cellular and is orchestrated by Th2 cells/cytokines. Recent studies are elucidating a role for additional cytokines such as IL-21 and IL-33 (Humphreys *et al.* 2008) and regulator cells, including the CD4⁺/Foxp3⁺ natural regulatory T cells and, alternatively activated macrophages (AAM) (Baumgart *et al.* 2006; Taylor *et al.* 2006; Persaud *et al.* 2007). While all immune cells can be considered as immunomodulatory (i.e., they affect other immune and stromal cells in their vicinity), regulatory T cells and AAMs (of which there are multiple types for each; Belkaid and Oldenhove, 2008; Reyes and Terrazas, 2007) have emerged as prototypic immunomodulators whose primary function is to switch-off or dampen immune reactions, or engage reactions that promote tissue recovery and restitution. Much research remains to be done to precisely define the biology of regulatory T cells and AAMs; however, the mobilization of both classes of cells following an infection with helminths would generate an immunoregulatory/immunosuppressive environment in which a spectrum of immune events would be inhibited. Thus, by extrapolation, the ability of products from the helminth to elicit regulatory T cell or AAM activity could contribute to worm infectivity and survival.

HELMINTH-DERIVED IMMUNOMODULATORS

An analysis of the biology of a parasite should include studying the mechanisms of the interaction of the parasite with the host immune system, so that potential parasite-derived immunomodulators (used to develop drugs to treat immuno-pathological

changes/disease) can be identified. During the last 40 years, a staggering number of helminth-derived immunomodulators have been partially characterized, both in terms of structure and bioactivity (we define an 'immunomodulator' as any molecule that regulates host immunity, but not antigens recognized by the host that are used to generate immune responses aimed at the destruction or elimination of the parasite from the host). Here, we focus on selected examples from the 3 main groups of parasitic helminths and draw the reader's attention to the fact that a literature search will reveal many other examples, all worthy of discussion, and excluded only on the basis of space restrictions (the examples given below are supplemented by additional material in Tables 1–3).

Trematodes

Schistosoma spp. Schistosomes, because of their impact on human health (King and Dangerfield-Cha, 2008), have been the focus of intense research efforts, and a variety of glycoconjugates have been extracted from the eggs and adult tegument (Bennett, 1963). Soluble egg antigen (SEA) from *S. mansoni* has been implicated in the host-response to infection, and has been shown consistently to have immunomodulatory properties. For example, SEA appears to be responsible for the reduction in CD4⁺ lymphocytes 6 weeks following infection, an event that was attributed to enhanced Fas-FasL-induced apoptosis involving CD19⁺ B cells (Lundy *et al.* 2001). Excretory/secretory (ES) products from schistosomes contain a 23 kDa molecule that results in T cell depletion *in vivo* via Fas-FasL killing. Exposure to SEA results in reduced levels of nitric oxide (NO) and T helper cell-type 1 (TH1) chemokines (CCL3) and cytokines (tumour necrosis factor- α (TNF α), IL-12) and an increased expression of markers indicative of AAMs. These events are all compatible with the mobilization of TH2-type cytokines (e.g. IL-4, IL-13) which is stereotypic of the mammalian immune response to infection by helminth parasites. The induction of AAMs is of particular note, as these cells exert anti-inflammatory effects and are active in tissue restitution after injury (Gordon and Taylor, 2005). Thus, the ability to use a constituent of SEA to induce this cellular response pharmacologically could be of direct health benefit.

Mice treated with SEA display endothelial cell proliferation and capillary formation which appear to be driven by a heat- and protease-insensitive molecule, indicative of a glycan (Kanse *et al.* 2005). These findings complement the work of van de Vijver *et al.* (2006) that detected LDN (GalNAc β 1,4GlcNAc) and LN (Gal β 1,4GlcNAc) modified glycans in SEA-stimulated granulomata. SEA has been shown to induce vascular endothelial growth

factor (VEGF) secretion (Loeffler *et al.* 2002) and fibroblast-stimulating factor (FSF-1) (Greenwel *et al.* 1993), which could contribute to the formation of the fibrous capsule around the granulomata, although it is unclear whether the stimulation of the production of either growth factor was due to a glycan component of the SEA. Nevertheless, one could speculate that SEA evoked growth factor release from host cells to stimulate angiogenesis could be used to enhance the recovery from tissue damage, although the potential to elicit metastatic events should not be overlooked.

In addition, while the exposure to SEA decreased the expression of Toll-like receptors (TLR) on macrophages, the responses of these cells to TLR agonists (e.g., lipopolysaccharide (LPS)) were not appreciably retarded (Joshi *et al.* 2008). These studies are intriguing, as they demonstrate a cross-talk between infection with parasitic helminths and responsiveness to bacteria and may illuminate a way in which one could exploit one type of infection to treat another. In addition, the complexity of the host response to infection, which is linked temporally or spatially and/or parasite-specific, is aptly illustrated by these studies.

The identification of a >100 kDa heat-insensitive, carbohydrate-positive fraction from *S. mansoni* (see Hayunga *et al.* 1979) was followed by studies characterizing the glycoproteins of schistosomes and notably led to the elucidation of lacto-N-fucopentose (LNFPIII) (carbohydrate containing the mammalian Lewis^X trisaccharide [Gal β 1,4[Fuca1,3]GlcNAc]) and lacto-N-neotetrose (LNT) (Ko *et al.* 1990). Analysis of LNT revealed that its injection into mice caused an expansion of a Gr1⁺ cell population which suppressed CD4⁺ T cell proliferation and cytokine production *via* an apoptosis-independent but cell contact-dependent mechanism (Terrazas *et al.* 2001). The authors postulated that this treatment 'imprinted' the Gr1⁺ cells with a TH2-type phenotype.

Since the Lewis^X antigen is present in the lymphocyte function-associated molecule-1 (LFA-1), the role of LNFPIII in lymphocyte proliferation was assessed and shown to stimulate IL-10 and prostaglandin E₂ (PGE₂) production from B-cells of infected mice without hampering antigen presentation (Velupillai and Harn, 1994). Also, the coculture of LNFPIII-treated macrophages and NK cells enhanced NK cell activation as gauged by the expression of CD69 and interferon- γ (IFN γ) production (Atochina and Harn, 2005). Lacto-N-fucopentose III also generates anti-inflammatory responses by signalling through Toll-like receptor 4 (TLR4) to induce the phosphorylation and activation of the mitogen-activated protein kinase (MAPK), ERK (Thomas *et al.* 2003).

The presence of fucose in the Lewis^X moiety of LNFPIII is essential for bioactivity (Okano *et al.*

Table 1. Selected examples of immunomodulatory molecules from trematodes

Species	Worm product	Bioactivity	Reference
<i>Schistosoma japonicum</i>	900 kDa ECF-SjE from homogenized eggs; pronase and heat insensitive glycoprotein; destroyed by periodate oxidation	<i>In vitro</i> eosinophil chemotaxis	Owhashi and Ishii (1982)
	440 kDa JAE-H and <440 kDa JAE-L glycoproteins from adult ES products	Eosinophil chemotactic factors; JAE-L also induces neutrophil chemotaxis	Horii <i>et al.</i> (1984)
<i>Schistosoma mansoni</i>	Analogues of adrenocorticotrophin (ACTH) and α -melanotrophin (α -MSH)	ACTH converts to α -MSH by polymorphonuclear cells via neutral endopeptidase; α -MSH inhibits leukocyte adherence and immunosuppressive	Duvaux-Miret <i>et al.</i> (1992)
	Various surface glycans	Fucose and Galactose linked to bovine serum albumin reduced ERK and PKC phosphorylation and phagocytosis in <i>Lymnaea stagnalis</i> haemocytes	Plows <i>et al.</i> (2005)
<i>Fasciola hepatica</i>	25 kDa glutathione S-transferase in ES products	Detoxification of peroxides involved in oxidative stress	Guillou <i>et al.</i> (2007)
	ES products contain metal ion dependent glycosidases (β -galactosidase, β -N-acetylhexosaminidase and β -glucosidase)	May degrade host mucins rich in galactose, N-acetylglucosamine and N-acetylgalactosamine	Irwin <i>et al.</i> (2004)
	ES products	Prevent superoxide production by PMA-activated sheep and human neutrophils <i>in vitro</i> (a heat resistant ES component from the related species, <i>F. gigantica</i> , does the same)	Jefferies <i>et al.</i> (1997) El-Ghaysh <i>et al.</i> (1999)
<i>Diplostomum pseudopathaceum</i>	22–24 kDa lectin with homology to β 1,3 glucan binding protein localizes to the cercarial penetration glands	Agglutinates murine red blood cells and may facilitate tissue recognition and penetration	Mikes and Horak (2001)
<i>Paragonimus westermani</i>	27 kDa cysteine protease in ES products	Induces superoxide production and human eosinophil degranulation	Chung <i>et al.</i> (2008)
<i>Opisthorchis viverrini</i>	24 kDa thioredoxin peroxidase (TPx) isolated by genomic probing; also exists in ES products	Protects worm from reactive oxygen metabolites and may have other roles similar to <i>F. hepatica</i> TPx	Suttiprapa <i>et al.</i> (2008)

1999), since lacto-N-neotetrose lacking Lewis^X had no effect on DC Φ differentiation to an IL-4 producing phenotype in response to LPS. Importantly, the induction of accessory molecules is required to increase the strength of an immune response; DC-SIGN has strong affinity for intercellular adhesion molecule-3 (ICAM3) via the Lewis^X structure (Bogoevska *et al.* 2007), suggesting a possible mechanism used by LNFPIII to bind to DC Φ and interfere with cell-to-cell communication with other immune cells (Appelmek *et al.* 2003). The multivalency of LNFPIII is important for its bioactivity, since monovalent LNFPIII was shown to be incapable of inhibiting antigen-induced peripheral blood mononuclear cell (PBMC) activation (Velupillai *et al.* 2000). At the signalling level, LNFPIII increases nuclear factor (NF)- κ B activity in DC Φ to a level similar to LPS; however, a second phase (prolonged NF- κ B activation associated with I κ B β degradation) was not observed. Use of the nuclear import inhibitor, SN50, and NF κ B-deficient mice revealed that LNFPIII effects on DC Φ were indeed NF κ B-dependent (Thomas *et al.* 2005). Collectively, these studies suggest that LNFPIII (or possibly fragments thereof) could be used to extrinsically manipulate DC Φ function as a way of limiting immunopathological reactions/changes or controlling immune cell responsiveness.

The stimulation of IL-10 and IL-12p40 from murine peritoneal macrophages by cercarial ES products required the presence of TLR4, whereas the production of IL-6 was TLR4 independent (Jenkins *et al.* 2005). Furthermore, IL-6 release was reduced following protease treatment of the ES products suggesting a proteinaceous ligand, and sodium periodate ablated activity of the cercarial extract, indicating the presence of glycans. It is plausible that cercarial products contain proteins/glycoproteins that can act *via* TLR4, highlighting the promiscuity of TLR4 as more than an LPS receptor. Other TLRs have been implicated as receptors for schistosome ligands. Lyso-phosphatidylserine from *S. mansoni* induces DC Φ maturation and decreases IL-12p70 production *via* TLR2. The activity is abrogated by TLR2-blocking antibodies (van der Kleij *et al.* 2002), whereas host phosphatidylserine neither activates TLR2 nor induces DC Φ maturation. Additionally, phosphoserine alone is not effective in this system, indicating that the acyl and phosphoserine groups of *S. mansoni* lysophosphatidylserine are required for the effect. Inhibition of lysophosphatidylserine acyltransferase (LPC-AT) did not affect TLR4 expression or the binding of LPS, but it did prevent TLR4 translocation to lipid rafts and thus blocked the formation of the TLR4 complex with a resultant decrease in TNF α and IL-6 production (Jackson *et al.* 2008). It is possible that *S. mansoni* lysophosphatidylserine also signals through TLR4 in a similar manner. Studies such

as these illustrate an emergent theme, in which responses to helminth-derived molecules occur *via* receptors that have been defined on the basis of their ability to recognize conserved microbe-derived products; however, the data must be interpreted with caution, since the possibility of bacterial contamination of the helminth extracts is omnipresent.

Fasciola hepatica. ES products from adult *F. hepatica* (FhES) contain a glycan that is cytotoxic to eosinophils and acts *via* tyrosine kinase and caspase-dependent mechanisms (Serradell *et al.* 2007). Induction of AAMs often accompanies an infection with helminths, and it has been reported that *F. hepatica* thioredoxin peroxidase elicits AAMs production (Donnelly *et al.* 2005). Numerous studies have shown that AAMs are induced by IL-4/IL-13 (Stein *et al.* 1992); however, in the case of *F. hepatica* thioredoxin peroxidase, this conversion appeared to be IL-4-independent (Donnelly *et al.* 2005). The peroxidase also increased transforming growth factor- β (TGF β) and PGE₂ while decreasing IL-10, IL-12 and IL-18 from stimulated macrophages – such events *in vivo* would skew the immune response in favour of immunoregulation/immunosuppression and away from TH1-dominated pro-inflammatory reactions. Complementing these findings, the *in vitro* treatment of bovine macrophages with FhES products inhibited IFN γ and NO production in response to bacterial products and skewed the macrophage population towards an AAM phenotype; the latter event was lost by heat treatment of the ES products (Flynn and Mulcahy, 2008). This emphasizes that we must constantly be vigilant of the possibility that parasite-derived molecules, as well as those produced by the host (e.g., cytokines), can modify the host's immunity. Finally, cathepsin L appears to be a dominant molecule in FhES and may aid *F. hepatica* by degrading host tissue to assist parasite entry/migration, and interfering with the host's immune response by, for example, the degradation of IgG (Smith *et al.* 1993; O'Neill *et al.* 2001).

Cestodes

Pseudophyllidean and cyclophyllidean tapeworms usually elicit limited damage in the alimentary tract of their definitive host. However, they are recognized by their host (i.e., anti-worm antibodies are generated) so, the ability to suppress or negate host immunity would remain a priority for the helminth (see Table 2).

Taenia crassiceps. T cells from *T. crassiceps*-infected mice are hypo-proliferative and, in the presence of excretory/secretory (TcES) products, show diminished responses to mitogen stimulation *in vitro* (Spolski *et al.* 2000, Scuitto *et al.* 1995). Administration of TcES products to mice caused expansion of F4/80⁺ Gr1⁺ cells, and these cells

Table 2. Selected examples of immunomodulatory molecules from cestodes

Cestode	Worm product	Bioactivity	Reference
<i>Echinococcus multilocularis</i>	ES product	Modifies macrophage accessory function leading to inhibition of lymphocyte proliferation <i>in vitro</i>	Rakha <i>et al.</i> (1991)
	Metacestode glycosphingolipids	Inhibits PBMC proliferation; decreases IL-2/IL-2 receptor expression	Persat <i>et al.</i> (1996)
	14-3-3 proteins in ES products	Inhibits LPS-induced nitric oxide production by rat macrophages	Andrade <i>et al.</i> (2004)
<i>Echinococcus granulosus</i>	E4 Gal α 1,4Gal glycan from protoscolices extract	Inhibits mitogenic-induced proliferation; increases IL-10	Dematteis <i>et al.</i> (2001)
	Antigen B protease inhibitor in hydatid cyst fluid	Inhibits neutrophil recruitment <i>in vitro</i> ; increased production of IL-4 and IL-13 by human PBMCs; impairs human DC Φ differentiation but polarizes maturation of immature DC Φ to TH2	Shepherd <i>et al.</i> (1991) Rigano <i>et al.</i> (2001, 2007)
<i>Taenia multiceps</i>	Glucose lectin in cyst fluid	May have similar function as ES product of <i>E. multilocularis</i>	Judson <i>et al.</i> (1987)
<i>Taenia taeniaeformis</i>	19.5 kDa proteinase inhibitor from somatic extract	Inhibits splenocyte proliferation in response to mitogenic and antigenic stimuli	Leid <i>et al.</i> (1984, 1986) Suquet <i>et al.</i> (1984)
	ES products	Sequesters complement; decreases mitogen-induced splenocyte proliferation IL-2 production	Rikishia <i>et al.</i> (1985) Burger <i>et al.</i> (1986)
<i>Taenia crassiceps</i>	Interferon- γ analogue (p66) in ES products	Enhances mitogen-induced splenocyte proliferation and IFN γ and IL-10 production; increases nitric oxide production by macrophages increases	Spolski <i>et al.</i> (2002)
<i>Taenia solium</i>	RNA peptide in metacestode extracts	Inhibits response to <i>Salmonella</i> antigens as well as mitogen-induced proliferation and inflammatory responses to metacestode antigens, correlating with decreased IL-2, IFN γ and IL-4; reduced TNF α production by macrophages following LPS stimulation	Arechavaleta <i>et al.</i> (1998)

were capable of blocking T cell activation *in vivo*. Glycans in the TcES were essential for this response, since the treatment with sodium periodate significantly reduced F4/80⁺ Gr1⁺ cells in the peritoneal cavity of infected mice and the production of cytokines and AAM (Gomez-Garcia *et al.* 2005). Extracts from the adult worm also directed a strong TH2 polarization (IL-4 production) and enhanced antibody production (IgG₁ and IgE) in mice co-treated with ovalbumin (Gomez-Garcia *et al.* 2006). Concomitant with these events was an increased expression of markers indicative of AAMs, a predicted decrease in IFN γ and iNOS expression and increased synthesis of IL-10 and TGF β . Like the ES products, the glycan component was critical for these effects of a crude antigen preparation from the adult worm, because all were abolished or significantly inhibited by sodium periodate treatment of the extract.

Extracts from the *T. crassiceps* metacestodes are highly mannosylated and contain both non-fucosylated and fucosylated structures, including the Lewis^X antigen (Dissanayake *et al.* 2004). In contrast to the adult worm antigen, the effect of which was unaltered by the knock-out of TLR4 (Gomez-Garcia *et al.* 2005), the ability of the glycans in the metacestode extract to elicit IL-6 production from murine peritoneal macrophages was reduced in cells lacking TLR2 and TLR4 (Dissanayake *et al.* 2004). The same researchers recently found that the terminal beta-(1-4)-galactose residue in the Lewis^X analogue can evoke IFN γ production by murine splenocytes and activate the NF- κ B pathway. The IFN γ production was reduced by inclusion of anti-TLR2, -4 and -6 antibodies in the assay (Dissanayake and Shahin, 2007), again reiterating the notion that helminth-derived products can exert their effects *via* TLRs.

Spirometra erinaceieuropaei. Excretory/secretory (SeES) products contain immunoglobulinases and a heat-sensitive factor that inhibits LPS- and lipoteichoic acid (LTA)-induced TNF α and IL-1 β production from murine macrophages (Diragahayu *et al.* 2004). The inhibitory effect of the SeES products was not due to the stimulation of PGE₂, IL-10 or TGF β in the target cells (Fukushima *et al.* 1993). The inhibition of macrophage production of TNF α , IL- β and NO was also shown in a study by Kina *et al.* (2005), in which a 90 kDa trypsin-sensitive glycoprotein was identified as the active component in the SeES products. Similarly, a 94 kDa component of SeES products could block LPS-induced MAPK (i.e., ERK1, ERK2 and p38MAPK) phosphorylation and the accompanying synthesis of TNF α (Diragahayu *et al.* 2002).

Hymenolepis diminuta. Extracts from adult *H. diminuta* reduced ConA-induced proliferation, IL-2

and IL-4 synthesis by murine spleen cells and human PBMC cells in a concentration-dependent manner (Wang and McKay, 2005). These events were mirrored, to a large extent, by use of ES products from the adult worm. A crude phosphate-buffered saline (PBS)-soluble extract of the adult worm stimulated IL-10 release from murine splenocytes and also contained a component with some structural similarity to the p40 chain of IL-12, as detected by enzyme-linked immunosorbent assay (ELISA) and immunoblotting. The bioactivity in both the worm extract and the ES products was within a >50 kDa fraction and was partially sensitive to boiling (Wang and McKay, 2005).

Nematodes

Parasitic nematodes have been the focus of extensive research efforts because of their major socio-economic impacts globally (Hall *et al.* 2008). Therefore, a number of immunomodulatory molecules have been identified from this phylum (Table 3). We focus this discussion by highlighting known molecules from *Nippostrongylus brasiliensis*, *Trichinella spiralis*, *Necator americanus*, *Trichuris suis*, *Ascaris suum* and *Acanthocheilonema viteae*.

Nippostrongylus brasiliensis. ES products from this parasite direct a range of immunomodulatory reactions. As examples, (i) they reduce LPS-induced IL-12p40 production and expression of a panel of cytokine/chemokine genes in murine bone-marrow-derived DC Φ , (ii) an undefined protein component induces IL-4, IL-10 and IgE production, with the IL-4 response requiring the presence of major histocompatibility class II (MHC II) antigens and so presumably MHC II-T cell restricted events, and (iii) suppress TH1-events via an IL-10-independent mechanism (Holland *et al.* 2000, 2005; Balic *et al.* 2004). The characterization of the ES products revealed the presence of a serine acetylhydrolase capable of reducing intestinal anaphylactic responses by inactivating platelet activating factor and blocking neutrophil and eosinophil aggregation (Blackburn and Selkirk, 1992). The value of inhibiting eosinophil activity to the helminth is intuitive, and the use of this acetylhydrolase would be beneficial in limiting the pro-inflammatory effects of granulated phagocytes in other diseases. Furthermore, NbES blocked goblet cell hyperplasia, eosinophilia, neutrophilia and airway hyper-responsiveness in murine models of asthma (Keir *et al.* 2004; Trujillo-Vargas *et al.* 2007). The active component/s was/were heat and protease insensitive, suggesting a glycan-type molecule. When NbES was administered intraperitoneally to ovalbumin-sensitized mice, they developed less pathology on ovalbumin re-challenge than untreated controls; in contrast, intranasal NbES induced significant lung inflammation (Marsland

Table 3. Selected examples of immunomodulatory molecules from nematodes

Nematode	Worm product	Bioactivity	Reference
<i>Ascaris suum</i>	Pseudocoelomic fluid protein(s)	Induces eosinophil and neutrophil chemotaxis; reduces bronchoalveolar lavage fluid cell infiltrate, IL-5 and IL-13 as well as expression of CD40 and CD86 and is therapeutic in murine asthma models	Tanaka <i>et al.</i> (1979) McConchie <i>et al.</i> (2006)
	Somatic extract from homogenized adults	Inhibits zymosan-induced murine arthritis; reduces hypersensitivity to ovalbumin independently of IL-10, and reduces cell proliferation, IL-2, IFN γ , eosinophil peroxidase, IL-4, IL-5 and eotaxin levels;	Rocha <i>et al.</i> (2008) Souza <i>et al.</i> (2002)
	200 kDa PAS-1 from adult somatic extract	Inhibits airway allergy induced by the worm allergen via inhibition macrophage cytokine release	Lima <i>et al.</i> (2002) Itami <i>et al.</i> (2005)
	Phosphorylcholine-conjugated glycosphingolipids from adults	Increases TNF α , IL-1 β and IL-6 from human PBMCs; decreases B cell proliferation; is pro-apoptotic; decreases TH1 cytokine production from LPS + IFN γ treated macrophages	Deehan <i>et al.</i> (2002) Kean <i>et al.</i> (2006) Katz <i>et al.</i> (2001)
<i>Anisakis simplex</i>	Heat sensitive larval homogenate	Eosinophil chemotaxis	Tanaka and Torisu (1978)
<i>Brugia malay</i>	47 kDa serine protease inhibitor in ES products	May inhibit neutrophil cathepsin G and elastase (disputed by Stanley and Stein (2003))	Zang <i>et al.</i> (1999)
	62 kDa asparaginyl-tRNA synthetase in ES products	Induces ERK1/2 MAPK phosphorylation via a G-protein coupled receptor; chemotactic for immature DC Φ , lymphocytes, eosinophils, neutrophils and cells positive for chemokine receptors CXCR1 and CXCR2,	Kron <i>et al.</i> (1995) Ramirez <i>et al.</i> (2006)
	Galectin-1 like molecule in ES products 12 kDa macrophage migration inhibitory factor (MIF) in ES products	Binds galactose containing glycoconjugates and may protect helminth from eosinophil and neutrophil mediated damage Likely to have similar function as <i>T. spiralis</i> MIF	Hewitson <i>et al.</i> (2008) Pastrana <i>et al.</i> (1998)
<i>Necator americanus</i>	ES products contain calreticulin	Identical to <i>H. contortus</i> calreticulin; C1q complement is bound which protects the nematode from surface damage while promoting feeding by sequestering pro-coagulants. May also bind C-reactive protein.	Kasper <i>et al.</i> (2001)
<i>Trichinella spiralis</i>	Encysting stage produces a 12 kDa macrophage migration inhibitory factor (MIF)-like molecule	Potential to regulate host macrophage responses	Wu <i>et al.</i> (2003)
<i>Toxocara canis</i>	O-glycans in ES products	May act as antibody decoys or to alter viscosity of host mucins	Gems and Maizels (1996) Loukas <i>et al.</i> (2000b)
	Tumour associated antigen ((GalNAc-O-Ser/Thr) (Tn)) and GalNAc polypeptide N-acetylgalactosaminyltransferase (ppGalNTase) ES products	The antigen is synthesized by the enzyme via abnormal elongation of O-glycans and interacts with macrophage C-type lectins and affects T cell activation Impair eosinophil-mediated damage through unknown mechanism	Clausen and Bennett (1996) Casaravilla <i>et al.</i> (2003) Giacomin <i>et al.</i> (2008)
<i>Teladorsagia circumcincta</i>	Adult and L4 ES products contain thioredoxin peroxidase (TPx)	Likely to have similar function to <i>F. hepatica</i> TPx in affecting TH1/TH2 cytokine balance	Craig <i>et al.</i> (2006)

<i>Dictyocaulus viviparus</i>	Possesses immunogenic N-glycan conjugated to phosphorylcholine as well as platelet activating factor	The latter may induce anti-PAF antibodies which may protect against eosinophil-mediated damage	Kooyman <i>et al.</i> (2007)
<i>Haemonchus contortus</i>	66 kDa glycoprotein in ES products 55 kDa glycoprotein in ES products 62 kDa calreticulin in ES products	Inhibits monocyte function Identical function as <i>A. caninum</i> 41 kDa product Prevents erythrocyte lysis and binds pro-coagulant Factor X, Ca ²⁺ and p24 which enables blood feeding by the worm; p24 has homology with C-reactive protein which is enhanced during helminth infection and induces tegumental damage via complement	Rathore <i>et al.</i> (2006) Anbu and Joshi (2008) Taylor and Hoole (1997) Kasper <i>et al.</i> (2001) Suchitra and Joshi (2005) Suchitra <i>et al.</i> (2008)
<i>Heligmosomoides polygyrus</i>	ES products	Similar anti-asthma activity as <i>N. brasiliensis</i> extracts except in this case the effects are mediated by IL-10 from CD4 ⁺ CD25 ⁺ Foxp3 ⁻ cells	Segura <i>et al.</i> (2007)
<i>Ancylostoma cantinum</i>	41 kDa glycoprotein in ES products	Inhibits neutrophil adhesion and blunts hydrogen peroxide production via CD11b/CD18 integrin	Moyle <i>et al.</i> (1994)

et al. 2005). Finally, and intriguingly, *N. brasiliensis* may be worthy of consideration in cancer therapy. When carcinogenic cells were injected with *N. brasiliensis* extracts, eosinophils were recruited and attacked the cells (i.e., as if they were a helminth), resulting in reduced size of the carcinoma (Frenoy, 2005).

Trichinella spiralis. Structural and biochemical analysis of *T. spiralis* ES products (*TrsES*) from larvae or adult worms revealed the presence of a nucleoside 5'-diphosphate phosphohydrolase (apyrase, 17 kDa), a 5' nucleotidase that preferentially targets AMP to generate adenosine and an adenosine deaminase (Gounaris, 2002; Gounaris *et al.* 2004). The ability to regulate AMP metabolism has a number of potential immunomodulatory effects, since adenosine is known to inhibit platelet aggregation, monocytes/macrophage TH1 cytokine production, and neutrophil synthesis of reactive oxygen species (Liaudet *et al.* 2002; Antonioli *et al.* 2008), and is protective in a murine model of colitis (Mabley *et al.* 2003). Also, adenosine stimulates epithelial electrogenic Cl⁻ secretion, which creates a driving force for water to move into the gut (Kottgen *et al.* 2003) and could act to flush *Trichinella* larvae as well as intestinal bacteria and protozoan pathogens from the host.

Necator americanus. Nematode-derived proteases serve a number of vital roles in larval exsheathment, the digestion of host tissue (facilitating invasion and migration) and in immune evasion (Gamble *et al.* 1989, McKerrow *et al.* 1990). *Necator americanus* ES (*NaES*) contains at least 7 cysteine proteases, 1 serine protease and various metalloproteases, some of which act as immunoglobulinases (Kumar and Pritchard, 1992). These enzymes exert a range of bioactivities, the majority of which are unknown. However, one can speculate that the cleavage of immunoglobulins (Igs) could result in reduced complement activation and that the activity of cells that bind Igs and are activated by antibody-cross-linkage by antigen (e.g., mast cells and eosinophils) would be reduced. Eosinophil recruitment *in vivo* could also be perturbed by, for example, the degradation of the chemokine, eotaxin-1 (Culley *et al.* 2000). Additionally, a *Necator americanus* protein was shown to evoke release of IFN γ from murine and human NK cells but not NK T cells (Hsieh *et al.* 2004).

Trichuris suis and *Ascaris suum*. *Trichuris suis* ES (*TrsES*) products contain heat-stable and protease-insensitive components which reduce the growth of the bacteria *Campylobacter jejuni*, *Escherichia coli* and *Staphylococcus aureus* (see Abner *et al.* 2001). High performance liquid chromatography (HPLC) did not reveal the presence of antibiotics (i.e., penicillin, streptomycin and amphotericin) in the

TrsES. The active component was <10 kDa in size. *A. suum* pseudocoelomic fluid (PCF) also contained a 6–14 kDa heat-stable, trypsin-sensitive molecule which could kill Gram-positive bacteria (e.g., *S. aureus*) (Wardlaw *et al.* 1994). The molecule was designated ASABF (*Ascaris suum* antibacterial factor) (Kato and Komatsu, 1996) and is similar to insect defensins which penetrate Gram-positive bacterial membranes. Genes encoding antibacterial cecropins had been characterized in *A. suum*, and chemically synthesized versions are active against numerous species of bacteria (Pillai *et al.* 2005). With increasing incidence of antibiotic-resistant bacteria, the putative usefulness of these nematode-derived molecules is apparent.

Acanthocheilonema viteae. Nematode-derived cystatins (cysteine protease inhibitors) have been repeatedly shown to regulate immune cell activity. Cystatins from *A. viteae* (as well as those of other filarioids, including *Onchocerca volvulus* and *Brugia pahangi*) inhibit T cell proliferation, interfere with antigen processing and presentation, and enhance IL-10 production from stimulated murine spleen cells (Osborne and Devaney, 1999; reviewed by Hartmann and Lucius, 2003). Recently an *A. viteae*-derived cystatin has been demonstrated to reduce inflammation in a murine model of airway allergy (Schnoeller *et al.* 2008). Onchocystatin (*Ov17*) from *O. volvulus* was specifically shown to suppress PBMC proliferation by decreasing expression of the co-stimulatory molecule B7.2 (CD86) (Schonemeyer *et al.* 2001), and infection with *O. volvulus* resulted in IL-10/TGF β -mediated immune hypo-responsiveness (Doetze *et al.* 2000). In that study, the source of the IL-10 and TGF β was most likely host cells, but the possibility of helminth-derived analogues of these cytokines should not be overlooked. For example, a TGF β -like molecule has been isolated from the filarial nematodes *Brugia malayi* and *Brugia pahangi* (Gomez-Escobar *et al.* 2000).

The best described helminth immunomodulator is an *A. viteae* ES 62 kDa glycoprotein (ES-62) product. The bioactivity of ES-62 is critically dependent on phosphorylcholine (PC) attached to an N-glycan (likely GlcNAc) *via* acetylglucosaminyl-transferase I, such that recombinant ES-62 lacking PC is generally non-functional (Goodridge *et al.* 2004; Houston and Harnett, 2004; Houston *et al.* 2008). Homologues of ES-62 exist in related nematodes (Stepek *et al.* 2002, 2004; Hewitson *et al.* 2008). Harnett *et al.* (2004) continue to explore the bioactivity of ES-62 in cell culture systems and murine models. Thus, ES-62 has been shown to enhance the production of IL-4 by murine splenocytes (Harnett *et al.* 1999) and decrease IFN γ , TNF α , IL-1 β , IL-12 and IL-18 from DC Φ stimulated with TLR ligands (Whelan *et al.* 2000; Goodridge *et al.* 2001, 2004). Complementing

the latter, macrophages from mice implanted with mini-osmotic pumps containing ES-62 were hypo-responsive to bacterial stimuli. ES-62 treatment in a murine model of arthritis resulted in significantly less inflammatory disease, which was accompanied by reduced IFN γ , TNF α and IL-6, and increased IL-10 synthesis by stimulated lymphocytes (McInnes *et al.* 2003). Significantly, IFN γ and T cell proliferation remained suppressed for up to 50 days after treatment whereas TNF α and IL-6 suppression required ongoing exposure, as did the inhibition of the production of collagen-specific antibodies. ES-62 also prevented B lymphocyte proliferation, and less IgG_{2a} would contribute to reduced auto-immune driven arthritis (Wilson *et al.* 2003).

ES-62 inhibits phosphorylation of immunoreceptor tyrosine-based activation motifs (ITAM) following B cell receptor ligation *via* the induction of the Src homology protein (SHP-1) tyrosine phosphatase. This effectively prevents the recruitment of crucial signalling molecules, such as protein kinase C (PKC) (Goodridge *et al.* 2005 *a*). A similar inhibition of PKC activity may relate to the ability of ES-62 to inhibit TH1 cytokine production (Deehan *et al.* 1997). In contrast, ES-62 inhibition of mast cell mediator release following the cross-linking of Fc ϵ RI did not require tyrosine phosphatase activity (Melendez *et al.* 2007). Instead, the increase in intracellular Ca²⁺ elicited by receptor cross-linking was affected and mobilization of phospholipase D and sphingosine kinase was disrupted (Goodridge *et al.* 2005 *a*). Fc ϵ RI-mediated NF- κ B activation and the consequent synthesis of pro-inflammatory cytokines were abrogated by ES-62 through the disruption of the p50 and p65 subunits of NF- κ B (Melendez *et al.* 2007). Given the pivotal role that mast cells play in immediate hypersensitivity and allergic reactions, a molecule, such as ES-62, could have many therapeutic applications.

The effect of ES-62 on mast cells appears to require TLR4 (Melendez *et al.* 2007), and this is in agreement with the examination of the effects of ES-62 on murine macrophages and DC Φ . This is not entirely unexpected, since PC can be a component on pathogen-associated molecular patterns (Goodridge *et al.* 2005 *b*). Thus, cells deficient in TLR4 or MyD88 (a key intracellular adaptor molecule) were unaffected by ES-62. The utilization of MyD88 may be critical, since cytokine-inducing stimuli that do not require this molecule, for example, the ligation of TLR3 or the accessory molecule CD40 in macrophages, are not affected by ES-62 treatment (Goodridge *et al.* 2005 *b*). Moreover, a natural mutation in the TIR domain in the TLR4 gene of C3H/HeJ mice renders these animals unresponsive to LPS, yet the ES-62 response remains, indicating that the helminth-derived molecule uses TLR4 in an unorthodox manner. It should be noted that the inhibitory effect of ES-62 can

be overridden, as observed in *Toxoplasma gondii*-infected mice, in which immunity relies on TH1 cytokines from a variety of cell types (Couper *et al.* 2005), or when T or B cells are challenged with two TH1-inducing antigens from *O. volvulus* (see Al-Riyami *et al.* 2008).

While the goal of this commentary is to illustrate the immunomodulatory effects of helminth-derived molecules and the potential of these molecules to serve as drugs or templates for drug design, it would be naïve to think that all parasite products would be beneficial. This point is intuitive and we will not belabour it here, but for the sake of balance in this discussion, we draw the reader's attention to a few of many examples in which helminth-derived products elicit pathological reactions (Hill *et al.* 1993; Lee *et al.* 2006; Kim *et al.* 2008).

We have presented selected examples of immunomodulation by parasitic helminth-derived molecules. Since the essence of any host-parasite relationship is specificity, it follows that nuances of immunomodulation will be parasite-, and indeed life-cycle stage-specific. However, some commonality emerges from these studies and, as generalizations, helminth-derived products mobilize a TH2 cytokine response (and ablate TH-1 events), stimulate the synthesis of immunomodulatory/immunosuppressive molecules (e.g., IL-10, TGF β , PGE₂), and some evoke eosinophil (and neutrophil) apoptosis. All of these events would ameliorate inflammatory disease and, as a pharmacopoeia for new drugs, the potential of the parasitic helminths should not be underplayed. Yet, much remains to be explored, to move the field from a series of intriguing observations *in vitro* and in rodents toward the development of drugs with a safety profile compatible with testing in clinical trials. At such a juncture, addressing issues relating to the *in vivo* stability and pharmacodynamics, and even the delivery methods of the helminth-derived molecule or drug, will be critical if new therapeutic modalities are to be achieved.

HELMINTH GLYCOBIOLOGY

Parasitic helminths as a group express or secrete/excrete a variety of molecules that have immunomodulatory properties. These molecules can be classical proteins or lipid-based, and many others are glycoproteins. We have opted to focus on parasite glycoproteins as molecules that could serve as models for the development of anti-inflammatory or immunoregulatory drugs, and will describe how such molecules can be isolated, characterized and might interact directly with mammalian cells.

Glycan types

An important type of protein post-translational modification is glycosylation, in which sugar residues

are added on to a core protein/sugar sequence, resulting in antennary structures of differing complexity in terms of carbohydrate variety, linkage, presence of sulphated groups and antennae numbers. N-linked glycans are attached to asparagine (Asn) *via* the tripeptide sequence Asn-X-Ser/Thr, where X represents any amino acid except proline. N-glycans containing LDN (GalNAc β 1,3GlcNAc) in antennary regions are considered 'complex' and are often capped by sialic acid or fucose *via* sialyl- and fucotransferases. O-linked glycans are attached to serine (Ser) or threonine (Thr) and, unlike N-glycans, do not share a common core sugar. O-glycan diversity is due to the presence of 8 possible core sugars. 'Hybrids' contain mannose and N-acetyllactosamine attached to a Man₃GlcNAc₂ chitobiose core.

Glycan isolation

Technological advances in the last 30 years have revolutionized glycoconjugate analysis, and considerable data can now be gleaned from minute amounts of material. Thus, metabolic radio-isotope labelling of carbohydrates using [³H], [¹²⁵I] and [¹⁴C], HPLC and fast performance liquid chromatography (FPLC), lectin affinity chromatography, followed by borate elution to increase the yield of glycans, coupled with more traditional approaches (e.g., high temperature [95 °C] hydrazinolysis, enzymatic cleavage and the use of periodate/borohydride to linearize sugar rings and remove reactive aldehydes) can yield considerable data on the nature of complex glycoproteins. Thus, mannose and fucose residues can be labelled using [2-³H]-mannose, sialic acid, GalNAc and GlcNAc with [6-³H]-glucosamine, whereas [6-³H]-galactose labels Gal and Glc. Further fractionation of labelled glycans can be accomplished *via* chromatography using agarose saturated with a single lectin type that reacts with carbohydrates in specific linkages (e.g., *Phaseolus vulgaris* agglutinin-agarose for tri- and tetra-antennary N-glycans possessing α -mannose substituted at C-2 and C-6 with Gal β 1,4GlcNAc) (Morelle and Michaelski, 2004). O-linked glycans have been analysed following β -elimination/release by mild alkaline and sodium borohydride treatment. However, this procedure releases some N-linked glycans and destroys O-acetyl groups, both of which can be avoided without lowering the yield of released glycans by using ammonia-borane β -elimination, with the added benefit that this treatment improves analysis *via* MALDI-MS (matrix assisted laser desorption/ionization time of flight) or CE-ESI-MS (capillary electrophoresis coupled to electrospray ionization mass spectrometry) (Huang *et al.* 2002).

All of these methodologies have been used to characterize helminth-derived glycans (Kang *et al.* 1993; Schallig and van Leeuwen, 1996; Casaravilla

et al. 2003). For example, PNGase F and A treatment or methanolysis-evoked release of radioisotope labelled N-glycans revealed fucosylated structures in the tegument of adult *S. mansoni* (see Nyame *et al.* 1989) and in H11 from *Haemonchus contortus* (see Haslam *et al.* 1996), while *Tetragonolobus purpureas* agglutinin-agarose electrophoresis and immunoblotting of metabolically labelled fractions with immune antisera demonstrated the Lewis^X antigen in schistosome glycoproteins (Srivatsan *et al.* 1992).

Other exquisitely sensitive techniques, such as fast atom bombardment (FAB)-MS, have provided structural information on glycans from *Dictyocaulus viviparus*, *H. contortus*, *T. spiralis*, *S. mansoni* and *O. volvulus* (see Haslam *et al.* 1996, 1999, 2000) as well as demonstrating O-glycan abundance in *Toxocara canis* ES products and in the glycocalyx of schistosome cercariae (Khoo *et al.* 1995). FAB-MS also demonstrated tyvelose (3,6 dideoxy-D-arabinohexose) on tri- and tetra-antennary glycans of *T. spiralis*, a modification previously regarded as unique to Gram-negative bacteria (Wisnewski *et al.* 1993). Although FAB-MS provides high sensitivity, MALDI-TOF mass spectrometry is ~10–100 times more sensitive (Huberty *et al.* 1993) and allowed the structure of the IL-4-inducing ES glycoprotein of *S. mansoni*, IPSEa1 to be characterized (Wuhrer *et al.* 2006).

Peptide mass finger-printing (PMF) determines protein identity following resolution by polyacrylamide electrophoresis. Resolved proteins are excised, cleaved by proteolysis, their masses determined by MALDI-TOF-MS and compared against a proteome database. When combined with lectin chromatography, mass spectrometry revealed the Lewis^Y antigen (Fuc α 1,2Gal β 1,4(Fuc α 1,3)GlcNAc β 1) as a modification of tegumental, but not ES glycoproteins from *S. bovis* (see Ramajo-Hernandez *et al.* 2007). Analysis by nanoESI (nano ElectroSpray Ionization) mass spectrometry enabled the characterization of the structures of immunogenic glycosphingolipids from *S. mansoni* (see Wuhrer *et al.* 2002, 2004). When nanoESI is combined with fourier transform ion cyclotron resonance (FTICR), analysis of intact, heavily sialylated glycoproteins can proceed without the need for chromatography, enzymatic treatment or derivitization (Nagy *et al.* 2004). FTICR can include fragmentation by electron capture dissociation (ECD), which cleaves protein backbones of glycoproteins without removing glycosylated regions, allowing glycosylation site determination. Combining ECD with external accumulation (XA) reduces 'background' while increasing resolution of unstable products, and was used to analyse <10 kDa glycoproteins containing sialic acids O-linked to GalNAc (Haselmann *et al.* 2001). Recently, quadrupole ion trap (QIT) technology has been coupled to ESI and MALDI-TOF

MS for glycoprotein analysis, in order to facilitate further molecular resolution of the isolated molecules (Demelbauer *et al.* 2004). When coupled to MALDI-TOF, nuclear magnetic resonance (NMR) radio-isotope labelling and glycosidase treatment, the structures of glycosphingolipids from *A. suum* were obtained (Friedl *et al.* 2003). The technology has culminated in quadrupole orthogonal acceleration time of flight mass spectrometry (Q-TOF) (Morris *et al.* 1997), which may be the most sensitive means for glycoprotein analysis.

These techniques can be applied to pure compounds, and many are suitable for analysis of mixtures of molecules. For example, CE-ESI-MS is superior to both CE and HPLC, and provides enhanced resolving power and separation of molecules based on electrophoretic mobility. Sheathless CE-ESI avoids sample dilution while increasing ionization and resolution efficacy. These qualities make CE-ESI-MS analysis attractive for glycoprotein investigations (Bateman *et al.* 1998). However, mass spectrometry has proven to be indispensable for biological analysis and has enabled comparisons of glycans among helminths (see Morelle and Michaelski, 2004). From this brief account, it is clear that the combination of these techniques with genomics, proteomics and glycomics have brought us to a point where accelerated progress in the elucidation of helminth-derived molecules (e.g., glycans) would be anticipated.

Are Siglecs receptors for helminth-derived molecules?

Some glycans from parasitic helminths appear to signal through TLRs. However, there are additional mammalian receptors which primarily recognize carbohydrates that are involved in immunomodulation, and these may have been 'co-opted' into the parasites strategy to overcome or skew host immunity. Siglec (sialic acid immunoglobulin-like lectin) receptors are expressed by leukocytes and recognize sialic acids; for example, Siglecs 1,3 and 4 recognize terminal α 2,3 sialic acid (Powell and Varki, 1994; Freeman *et al.* 1995; Collins *et al.* 1997); Siglec-2 binds α 2,6 sialic acid (Kelm *et al.* 1994); and, Siglecs 5, 7 and 9 recognize both sugar moieties (Cornish *et al.* 1998; Zhang *et al.* 2000). The cytoplasmic tails of Siglecs possess immunoreceptor tyrosine-based inhibitory motifs (ITIMs) and, so, Siglec activation is often associated with immunosuppression. For instance, the ligation of Siglec 9, and to a lesser extent Siglec 5, reduced TNF α production by macrophages while increasing IL-10 synthesis (Ando *et al.* 2008). Also, the repression of Siglec 3 (or CD33) expression with siRNA increased the spontaneous release of IL-1 β from monocytes (Lajaunias *et al.* 2005). These findings, and others (see Ikehara *et al.* 2004), lend credence to the hypothesis that Siglec-mediated

recognition of 'self' sialic acid by leukocytes may serve to generally dampen immune responses and perhaps promote tolerogenic events.

The ligation of Siglec 8 on eosinophils (and Siglec 9 on neutrophils; von Gunten *et al.* 2005) induces apoptosis (Nutku *et al.* 2003), and the ability of helminths to trigger such a response would be to their benefit. The published literature on parasitic helminths possessing sialic acid-based glycoproteins is conflicting, with many studies suggesting that helminths do not synthesize them, and there is an equally convincing body of evidence demonstrating that sialic acid reactivity can be detected in worm extracts or ES products from many helminth species (McDiarmid and Podesta, 1984; Apinhasmit *et al.* 2000; Elayoubi and Craig, 2004; Johnston, unpublished observations). We feel that this issue is deserving of clarity that will only come with additional well-controlled experimentation. Currently, there are no data on helminth-derived molecules that can bind Siglecs, and thus we can only speculate on the possibility of helminth-derived glycans-Siglec receptor interactions as a mechanism for helminth parasites to elicit a general immunosuppression in their hosts.

Are galectins receptors for helminth-derived molecules?

Beta-galactose binding lectins (galectins) are secreted carbohydrates that bind cell surface and matrix-associated glycans. Fifteen galectins have been identified in mammals and each possesses a conserved carbohydrate recognition domain (CRD) which binds N-acetylglucosamine (LN), Gal β 1,4GlcNAc or Gal β 1,3GlcNAc on N- and O-linked glycans (Milos and Zaliik, 1983). Galectins can be monovalent (one CRD), bivalent or chimeric (one CRD plus a unique amino terminal sequence). There are many examples of galectin modulation of immune activity, such as the induction of apoptosis, enhancement of macrophage phagocytosis, TH2-TH1 skewing and the induction of regulatory T cells (Rabinovich *et al.* 2007). Current data suggest that the helminth-derived glycans can interfere with galectin-driven events. Galectin-1 ligation by helminth galactose can potentially decrease TNF α and IFN γ and prevent mast cell degranulation and neutrophil activity (Rabinovich *et al.* 2002). The galectin-3 receptor recognizes lactose-containing glycans (LN and LDN) from *S. mansoni*. This interaction has been implicated in granuloma formation (van den Berg *et al.* 2004) (recently disputed by Bickle and Helmby, 2007) and in eosinophil-mediated cytotoxicity (Truong *et al.* 1993), although the effect on eosinophils could be reduced by the galectin-3 inhibition of IL-5 production (Cortegano *et al.* 1998). In addition, mice in which galectin-3 expression is reduced (*via* genetic manipulation)

display reduced numbers of splenic T and B cells, less liver pathology and increased levels of TH1-type cytokines following *S. mansoni* infection (Breuilh *et al.* 2007). Similarly, galectin-3 appears to provide a degree of protection against *Clonorchis sinensis*-induced cholangiocarcinoma (see Junking *et al.* 2008). With the exception of galectin-3 (Yang *et al.* 1996), galectin activation of a cell often induces apoptosis (reviewed by Hernandez and Baum, 2002), an event that can be inhibited by α 2,6 sialic acid (Leffler and Barondes, 1986). Since many helminths produce sialic acids, the release of these glycans from the worm could increase the life-span of immune cells.

Many helminths also secrete galectins (Turner *et al.* 2008). For example, a 34 kDa galectin in *O. volvulus* is temporally expressed by third-stage larvae (L3) and adults (Joesph *et al.* 2000), shares 40% homology to a *Teladorsagia circumcincta*- and *H. contortus*-derived galectin (Greenhalgh *et al.* 1999), and is recognized by sera from infected individuals. The related galectin from *O. volvulus*, OvGalBP, binds IgE and may protect against eosinophil and neutrophil-mediated damage of the helminth (Klion *et al.* 1994). Galectin-1 administration is therapeutic in a plethora of murine models of autoimmune disease, where it is associated with reduced TH1 and TH17 events and increased CD4⁺ T cell apoptosis (Salatino *et al.* 2008). A galectin-1 like molecule has been identified in nematodes (Table 3). Moreover, chitin (a GlcNAc polymer) is abundant in helminths and profoundly affects immunity by mobilizing AAMs, eosinophils and basophils *in vivo* (Reese *et al.* 2007).

Are C-type lectins receptors for helminth-derived molecules?

C-type lectins (CTLs) are Ca²⁺-dependent carbohydrate receptors involved in signalling, antigen presentation and cell-cell adhesions (Pyz *et al.* 2006). CTLs include the collectins, which bind mannan proteins, the selectins, which bind sialylated/sulphated glycans, and the macrophage mannose receptor (MMR). A few studies have shown that parasitic helminths produce lectin-type molecules and molecules that bind host lectins, thus imparting the potential to affect processes as diverse as macrophage activation, immune cell trafficking and dendritic cell activity. For instance, mannan-binding lectin (MBL) interacts with glycans on *S. mansoni* and *T. spiralis*, resulting in complement fixation (Gruden-Movsesijan *et al.* 2003). Similarly, the macrophage galactose-type lectin (MGL), which is also expressed on DC Φ , binds the Lewis^x antigen and α/β GalNAc, both of which occur in *S. mansoni* LDN and LDNF (van Vliet *et al.* 2005). Lectin-targeting of the worm surface would be aimed at worm destruction, although the shedding of the

glycan from the parasite could be a component of an immune evasion strategy. Parasite-secreted CTL-like products have been identified and include ES products of ~32 and 70 kDa from *Toxocara canis* which resemble the MMR (CD206) and the low affinity IgE receptor (CD23), respectively (Loukas *et al.* 1999, 2000a). Lectins encoded by ESTs in *Ancylostoma ceylanicum* and *Ascaris suum* exist in the GenBank database and a *Necator americanus* lectin is predicted from transcriptome analysis (Daub *et al.* 2000). Exactly how helminth-derived glycans or lectin-like molecules modulate mammalian immune responses is not known; however, there is potential for pure forms of these molecules to be used as potent drugs. For example, glycans and SEA from *S. mansoni* can bind to DC-SIGN (Meyer *et al.* 2005; van Liempt *et al.* 2007). DC-SIGN is important in recognizing mannose-containing carbohydrates and is also an activation signal that helps direct DC Φ migration from tissues to lymph nodes to initiate primary adaptive immune responses.

CONCLUSIONS

The concept of helminth-derived molecules as a source of immunomodulatory agents is fully established (Hartnett and Harnett, 2008), and the literature is flush with examples of how crude extracts and partially characterized molecules from many species of parasitic helminth can affect virtually all aspects of innate and adaptive immune responses. To date, this wealth of data has not been converted into a slew of new drugs. While the promise of parasitic helminths as a pharmacopeia has yet to be fulfilled, this should not deter future research in this area, as the potential pay-back in terms of new drugs is immense. Indeed, one could foresee that a number of novel drug candidates will be forthcoming in the next 5–10 years, given the advent of technology for the purification and characterization of molecules from small amounts of tissues and the synthesis of agents for testing in high throughput immunological assays. We subscribe to, and will leave the reader with the view presented by Falcone and Pritchard (2005): “The fact that the compounds from parasites might have evolved to unlock the human immune system selectively will also result in less attrition when compounds are tested for safety and adverse activity profiles”.

REFERENCES

- Abner, S. R., Parthasarathy, G., Hill, D. E. and Mansfield, L. S.** (2001). *Trichuris suis*: detection of antibacterial activity in excretory-secretory products from adults. *Experimental Parasitology* **99**, 26–36.
- Al-Riyami, L., Egan, C. A., Bradley, J. E., Lustigman, S. and Harnett, W.** (2008). Failure of ES-62 to inhibit T-helper type I responses to other filarial nematode antigens. *Parasite Immunology* **30**, 304–308.
- Anbu, K. A. and Joshi, P.** (2008). Identification of a 55 kDa *Haemonchus contortus* excretory/secretory glycoprotein as a neutrophil inhibitory factor. *Parasite Immunology* **30**, 23–30.
- Ando, M., Tu, W., Nishijima, K. I. and Iijima, S.** (2008). Siglec-9 enhances IL-10 production in macrophages via tyrosine based motifs. *Biochemical and Biophysical Research Communications* **369**, 878–883.
- Andrade, M. A., Siles-Lucas, M., Espinoza, E., Perez Arellano, J. L., Gottstein, B. and Muro, A.** (2004). *Echinococcus multilocularis* laminated-layer components and the E14t 14-3-3 recombinant protein decrease NO production by activated rat macrophages *in vitro*. *Nitric Oxide* **10**, 150–155.
- Antonoli, L., Fornai, M., Colucci, R., Ghisu, N., Tuccori, M., Del Tacca, M. and Blandizzi, C.** (2008). Pharmacological modulation of adenosine system: novel options for treatment of inflammatory bowel disease. *Inflammatory Bowel Disease* **14**, 566–574.
- Antony, R. M., Rutitzky, L. I., Urban, J. F., Jr., Stadecker, M. J. and Gause, W. C.** (2007). Protective immune mechanism in helminth infection. *Nature Reviews in Immunology* **7**, 975–987.
- Apinhasmit, W., Sobhon, P., Tarasub, C., Mothong, W., Saitongdee, P., Sretarugsa, P., Wanichanon, C. and Upatham, E. S.** (2000). *Opisthorchis viverrini*: ultrastructure and cytochemistry of the glycocalyx of the tegument. *Journal of Helminthology* **74**, 23–29.
- Appelmelk, B. J., Van Die, I., Van Vleit, S. J., Vandembroucke-Grauls, C. M., Geijtenbeek, T. B. and Van Kooyk, Y.** (2003). Cutting edge: carbohydrate profiling identifies new pathogens that interact with dendritic cell specific ICAM-3 grabbing nonintegrin on dendritic cells. *Journal of Immunology* **170**, 1635–1639.
- Arechavleta, F., Mollinari, J. L. and Tato, P.** (1998). A *Taenia solium* metacestode factor nonspecifically inhibits cytokine production. *Parasitology Research* **84**, 117–122.
- Atochina, O. and Harn, D.** (2005). LNFPIII/LeX-stimulated macrophages activate natural killer cells via CD40-CD40L interaction. *Clinical and Diagnostic Laboratory Immunology* **12**, 1041–1049.
- Balic, A., Harcus, Y., Holland, M. J. and Maizels, R. M.** (2004). Selective maturation of dendritic cells by *Nippostrongylus brasiliensis*-secreted proteins drives TH2 immune responses. *European Journal of Immunology* **34**, 3047–3059.
- Bateman, K. P., White, R. L. and Thibault, P.** (1998). Evaluation of adsorption preconcentration/capillary zone electrophoresis/nano-electrospray mass spectrometry for peptide and glycoprotein analyses. *Journal of Mass Spectrometry* **33**, 1109–1123.
- Baumgart, M., Tomkins, F., Leng, J. and Hess, M.** (2006). Naturally occurring CD4⁺Foxp3⁺ regulatory T cells are an essential, IL-10-independent part of the immunoregulatory network in *Schistosoma mansoni* egg-induced inflammation. *Journal of Immunology* **176**, 5374–5387.
- Belkaid, Y. and Oldenhove, G.** (2008). Tuning microenvironments: induction of regulatory T cells by dendritic cells. *Immunity* **29**, 362–371.
- Bennett, H.** (1963). Morphological aspects of extracellular polysaccharides. *Journal of Histochemistry and Cytochemistry* **11**, 14–23.

- Bickle, Q. and Helmsby, H.** (2007). Lack of galectin-3 involvement in murine intestinal nematode and schistosome infection. *Parasite Immunology* **29**, 93–100.
- Blackburn, C. C. and Selkirk, M. E.** (1992). Inactivation of platelet-activating factor by a putative acetylhydrolase from the gastrointestinal nematode parasite, *Nippostrongylus brasiliensis*. *Immunology* **75**, 41–46.
- Bogoevska, V., Nallau, P., Lucka, L., Grunow, D., Klampe, B., Uotila, L. M., Samsen, A., Gahmberg, C. G. and Wagener, C.** (2007). DC-SIGN binds ICAM-3 isolated from peripheral human leukocytes through Lewis X residues. *Glycobiology* **17**, 324–333.
- Breuilh, L., Vanhoutte, F., Fontaine, J., van Stijn, C. M. W., Tillie-LeBlond, I., Capron, M., Faveeuw, C., Jouault, T., van Die, I., Gosset, P. and Trottein, F.** (2007). Galectin-3 modulates autoimmune and inflammatory responses during helminthic infection: impact of galectin-3 deficiency on the functions of dendritic cells. *Infection and Immunity* **75**, 5148–5157.
- Burger, C. J., Rikihisa, Y. and Lin, Y. C.** (1986). *Taenia taeniformes*: inhibition of mitogen induced proliferation and interleukin-2 production in rat splenocytes by larval *in vitro* product. *Experimental Parasitology* **62**, 216–222.
- Carod-Artal, F. J.** (2008). Neurological complications of *Schistosoma* infection. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **102**, 107–116.
- Casaravilla, C., Friere, T., Malgor, R., Medeiros, A., Osinaga, E. and Carmona, C.** (2003). Mucin type O-glycosylation in helminth parasites from major taxonomic groups: evidence for widespread distribution of the Tn antigen (GalNAc-Ser/Thr) and identification of UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferase activity. *Journal of Parasitology* **89**, 709–715.
- Chen, C. C., Louie, S., McCormick, B., Walker, W. A. and Shi, H. N.** (2005). Concurrent infection with an intestinal helminth parasite impairs host resistance to enteric *Citrobacter rodentium* and enhances *Citrobacter*-induced colitis in mice. *Infection and Immunity* **73**, 5468–5481.
- Chung, Y. B., Kita, H. and Shin, M. H.** (2008). A 27 kDa cysteine protease secreted by newly excysted *Paragonimus westermani* metacercariae induces superoxide anion production and degranulation of human eosinophils. *Korean Journal of Parasitology* **46**, 95–99.
- Clausen, H. and Bennett, E.** (1996). A family of UDP-GalNAc: polypeptide N-acetylgalactosaminyltransferase controls the initiation of mucin type O-linked glycosylation. *Glycobiology* **6**, 635–646.
- Collins, B. E., Kiso, M., Hasegawa, A., Tropak, M. B., Roder, J. C., Crocker, P. R. and Schnaar, R. L.** (1997). Binding specificities of the sialoadhesion family of I-type lectins. Sialic acid linkage and substructure requirements for binding of myeloid associated glycoprotein, Schwann cell myelin protein and sialoadhesion. *Journal of Biological Chemistry* **272**, 16889–16895.
- Cornish, A. L., Freeman, S., Forbes, G., Ni, J., Zhang, M., Cepeda, M., Gentz, R., Augustus, M., Carter, K. C. and Crocker, P. R.** (1998). Characterization of Siglec-5, a novel glycoprotein expressed on myeloid cells related to CD33. *Blood* **92**, 2123–2132.
- Corteganao, I., del Pozo, V., Cardaba, B., de Andres, B., Gallardo, S., del Amo, A., Arrieta, I., Jurado, A., Palomino, P., Liu, F. T. and Lahoz, C.** (1998). Galectin-3 downregulates IL-5 gene expression on different cell types. *Journal of Immunology* **161**, 385–389.
- Couper, K. N., Chen, W., Houston, K. M., Harnett, W. and Johnson, L. L.** (2005). ES-62 is unable to modulate *Toxoplasma gondii*-driven TH1 responses and pathology. *Parasite Immunology* **27**, 147–150.
- Craig, H., Wastling, J. M. and Knox, D. P.** (2006). A preliminary proteomic survey of the *in vitro* excretory/secretory products of fourth stage larval and adult *Teladorsagia circumcincta*. *Parasitology* **132**, 535–543.
- Culley, F. J., Brown, A., Conroy, D. M., Sabroe, I., Pritchard, D. I. and Williams, T. J.** (2000). Eotaxin is specifically cleaved by hookworm metalloproteases preventing its action *in vitro* and *in vivo*. *Journal of Immunology* **165**, 6447–6453.
- Daub, J., Loukas, A., Pritchard, D. I. and Blaxter, M.** (2000). A survey of genes expressed in adults of the human hookworm, *Necator americanus*. *Parasitology* **120**, 171–184.
- Deehan, M. R., Goodridge, H. S., Blair, D., Lochnit, G., Dennis, R. D., Geyer, R., Harnett, M. M. and Harnett, W.** (2002). Immunomodulatory properties of *Ascaris suum* glycosphingolipids – phosphorylcholine and non-phosphorylcholine-dependent effects. *Parasite Immunology* **24**, 463–469.
- Deehan, M. R., Harnett, M. M. and Harnett, W.** (1997). A filarial nematode secreted product differentially modulates expression and activation of protein kinase C isoforms in B lymphocytes. *Journal of Immunology* **159**, 6105–6111.
- Dematteis, S., Piroto, F., Marques, J., Nieto, A., Orn, A. and Baz, A.** (2001). Modulation of the cellular immune response by a carbohydrate rich fraction from *Echinococcus granulosus* protoscolexes in infected or immunized Balb/c mice. *Parasite Immunology* **23**, 1–9.
- Demelbauer, U. M., Zehl, M., Plematl, A., Allmaier, G. and Rizzi, A.** (2004). Determination of glycopeptide structures by multistage mass spectrometry with low energy collision induced dissociation: comparison of electrospray ionization quadrupole ion trap and matrix assisted laser desorption ionization quadrupole ion trap reflection time of flight approaches. *Rapid Communications in Mass Spectrometry* **18**, 1575–1582.
- Diragahayu, P., Fukumoto, S., Tadamoto, S., Kina, Y. and Hirai, K.** (2004). Excretory/secretory products from plerocercoids of *Spirometra erinaceieuropaei* suppress interleukin-1 β gene expression in murine macrophages. *International Journal for Parasitology* **34**, 577–584.
- Diragahayu, P., Fukumoto, S., Miura, K. and Hirai, K.** (2002). Excretory/secretory products from plerocercoids of *Spirometra erinaceieuropaei* suppress the TNF α gene expression by reducing phosphorylation of ERK1/2 and p38 MAPK in macrophages. *International Journal for Parasitology* **32**, 1155–1162.
- Dissanayake, S., Amith, R. S. and Shahin, A.** (2004). *Taenia crassiceps* carbohydrates stimulate IL-6 expression in naive murine macrophages via Toll-like receptors (TLRs). *Molecular Immunology* **41**, 391–398.

- Dissanayake, S. and Shahin, A.** (2007). Induction of interferon- γ by *Taenia crassiceps* glycans and Lewis sugars in naive Balb/c spleen and peritoneal exudate cells. *Molecular Immunology* **44**, 1623–1630.
- Doetze, A., Satoguina, J., Buchard, G., Rau, T., Loliger, C., Flesicher, B. and Horeauf, A.** (2000). Antigen specific cellular hyporesponsiveness in a chronic human helminth infection is mediated by Th3/Tr1-type cytokines IL-10 and transforming growth factor β but not by a Th1 to Th2 shift. *Infection and Immunity* **12**, 623–630.
- Donnelly, S., O'Neill, S. M., Sekiya, M., Mulcahy, G. and Dalton, J. P.** (2005). Thioredoxin peroxidase secreted by *Fasciola hepatica* induces the alternative activation of macrophages. *Infection and Immunity* **73**, 166–173.
- Duvaux-Miret, O., Stefano, G. B., Smith, E. M., Dissous, C. and Capron, A.** (1992). Immunosuppression in the definitive and intermediate hosts of the human parasite *Schistosoma mansoni* by release of immunoreactive neuropeptides. *Proceedings of the National Academy of Sciences, USA* **89**, 778–781.
- Elayoubi, F. A. and Craig, P. S.** (2004). *Echinococcus granulosus* coproantigens: chromatographic fractionation and characterization. *Parasitology* **128**, 455–465.
- El-Gayash, A., Turner, R. J., Brophy, P. M. and Barrett, J.** (1999). Effect of *Fasciola gigantica* somatic extracts and excretory/secretory products on superoxide production by activated neutrophils. *Veterinary Parasitology* **84**, 91–100.
- Falcone, F. H. and Pritchard, D. I.** (2005). Parasite role reversal: worms on trial. *Trends in Parasitology* **21**, 157–160.
- Farah, I. O., Langoi, D., Nyaundi, J. and Hau, J.** (2007). Schistosoma-induced pathology is exacerbated and TH2 polarization is enhanced during pregnancy. *In Vivo* **21**, 599–602.
- Flynn, R. J. and Mulcahy, G.** (2008). Possible role for Toll-like receptors in interaction of *Fasciola hepatica* excretory-secretory products with bovine macrophages. *Infection and Immunity* **76**, 678–684.
- Freeman, S. D., Kelm, S., Barber, E. K. and Crocker, P. R.** (1995). Characterization of CD33 as a new member of the sialoadhesion family of cellular interaction molecules. *Blood* **85**, 2005–2012.
- Frenoy, G.** (2005). Methods for the reduction of malignant tumors by an eosinophil/helminth therapy. *Free Patents Online* <http://www.freepatentsonline.com/7335354.html>
- Friedl, C. H., Lochnit, G., Zahringer, U., Bahr, U. and Geyer, R.** (2003). Structural elucidation of zwitterionic carbohydrates derived from glycosphingolipids of the porcine parasitic nematode *Ascaris suum*. *The Biochemical Journal* **369**, 89–102.
- Fukushima, T., Isobe, A., Hojo, N., Shiwaku, K., Yamane, Y. and Torii, M.** (1993). The metabolism of arachidonic acid to prostaglandin E₂ in plerocercoids of *Spirometa erinacei*. *Parasitology Research* **79**, 634–638.
- Gamble, H. R., Purcell, J. P. and Fetterer, R. H.** (1989). Purification of a 44 kilodalton protease which mediates the ecdysis of infective *Haemonchus contortus* larvae. *Molecular and Biochemical Parasitology* **33**, 49–58.
- Gems, D. and Maizels, R. M.** (1996). An abundantly expressed mucin-like protein from *Toxocara canis* infective larvae: the precursor of the larval surface coat glycoproteins. *Proceedings of the National Academy of Sciences, USA* **93**, 1665–1670.
- Giacomin, P. R., Cava, M., Tumes, D. J., Gauld, A. D., Iddawela, D. R., McColl, S. R., Parsons, J. C., Gordon, D. L. and Dent, L. A.** (2008). *Toxocara canis* larval excretory/secretory proteins impair eosinophil dependent resistance of mice to *Nippostrongylus brasiliensis*. *Parasite Immunology* (epub ahead of print) (in the Press).
- Gomez-Escobar, N., Gregory, W. F. and Maizels, R. M.** (2000). Identification of *tgh-2*, a filarial nematode homolog of *Caenorhabditis elegans daf-7* and human transforming growth factor β , expressed in microfilarial and adult stages of *Brugia malayi*. *Infection and Immunity* **68**, 6402–6410.
- Gomez-Garcia, L., Lopez-Marin, L. M., Saaverdra, R., Reyes, J. L., Rodriguez-Sosa, M. and Terrazas, L. I.** (2005). Intact glycans from cestode antigens are involved in innate activation of myeloid suppressor cells. *Parasite Immunology* **27**, 395–405.
- Gomez-Garcia, L., Rivera-Montoya, I., Rodriguez-Sosa, M. and Terrazas, L. I.** (2006). Carbohydrate components of *Taenia crassiceps* metacestodes display TH2 adjuvant and anti-inflammatory properties when co-injected with bystander antigen. *Parasitology Research* **99**, 440–448.
- Goodridge, H. S., Deehan, M. R., Harnett, W. and Harnett, M. M.** (2005 a). Subversion of immunological signaling by a filarial nematode phosphorylcholine-containing secreted product. *Cellular Signaling* **17**, 11–16.
- Goodridge, H. S., Marshall, F. A., Else, K. J., Houston, K. M., Egan, C., Al-Riyami, L., Liew, F. Y., Harnett, W. and Harnett, M. M.** (2005 b). Immunomodulation via novel use of TLR4 by the filarial nematode phosphorylcholine-containing secreted product, ES-62. *Journal of Immunology* **174**, 284–293.
- Goodridge, H. S., Marshall, F. A., Wilson, E. H., Houston, K. M., Liew, F. Y., Harnett, M. M. and Harnett, W.** (2004). *In vivo* exposure of murine dendritic cell and macrophage bone marrow progenitors to the phosphorylcholine-containing filarial nematode glycoprotein ES-62 polarizes their differentiation to an anti-inflammatory phenotype. *Immunology* **113**, 491–498.
- Goodridge, H. S., Wilson, E. H., Harnett, W., Campbell, C. C., Harnett, M. M. and Liew, F. Y.** (2001). Modulation of macrophage cytokine production by ES-62, a secreted product of the filarial nematode *Acanthocheilonema vitae*. *Journal of Immunology* **167**, 940–945.
- Gordon, S. and Taylor, P. R.** (2005). Monocyte and macrophage heterogeneity. *Nature Reviews in Immunology* **5**, 953–964.
- Gounaris, K.** (2002). Nucleotidase cascades are catalysed by secreted proteins of the parasitic nematode *Trichinella spiralis*. *Infection and Immunity* **70**, 4917–4924.
- Gounaris, K., Selkirk, M. E. and Sadeghi, S. J.** (2004). A nucleotidase with unique catalytic properties is

- secreted by *Trichinella spiralis*. *Molecular and Biochemical Parasitology* **136**, 257–264.
- Greenhalgh, C. J., Beckham, S. A. and Newton, S. E.** (1999). Galectins from sheep gastrointestinal nematode parasites are highly conserved. *Molecular and Biochemical Parasitology* **98**, 285–289.
- Greenwel, P., Wyler, D. J., Rojkind, M. and Prakash, S.** (1993). Fibroblast stimulating factor 1, a novel lymphokine produced in schistosomal egg granulomas, stimulates liver fat storing cells *in vitro*. *Infection and Immunity* **61**, 3985–3987.
- Gruden-Movsesijan, A., Petrovic, M. and Sofronic-Milosavljevic, L.** (2003). Interaction of mannan-binding lectin with *Trichinella spiralis* glycoproteins, a possible innate immune mechanism. *Parasite Immunology* **25**, 545–552.
- Guillou, F., Roger, E., Mone, Y., Rognon, A., Grunau, C., Theron, A., Mitta, G., Coustau, C. and Gourbal, B. E.** (2007). Excretory-secretory proteome of larval *Schistosoma mansoni* and *Echinostoma caproni*, two parasites of *Biomphalaria glabrata*. *Molecular and Biochemical Parasitology* **155**, 45–56.
- Hall, A., Hewitt, G., Tuffrey, V. and de Silva, N.** (2008). A review and meta-analysis of the impact of intestinal worms on child growth and nutrition. *Maternal and Child Nutrition* **4** (Suppl.1), 118–136.
- Harnett, W., Deehan, M. D., Houston, K. M. and Harnett, M. M.** (1999). Immunomodulatory properties of a phosphorylcholine-containing secreted filarial glycoprotein. *Parasite Immunology* **21**, 601–608.
- Harnett, W., McInnes, I. B. and Harnett, M. H.** (2004). ES-62, a filarial nematode-derived immunomodulator with anti-inflammatory potential. *Immunology Letters* **94**, 27–33.
- Harnett, W. and Harnett, M. M.** (2008). Therapeutic immunomodulators from nematode parasites. *Expert Reviews in Molecular Medicine* **10**, 1–13.
- Hartmann, S. and Lucius, R.** (2003). Modulation of host immune responses by nematode cystatins. *International Journal for Parasitology* **33**, 1291–1302.
- Haselmann, K. F., Budnik, B. A., Olsen, J. V., Nielsen, M. L., Reis, C. A., Clausen, H., Johnsen, A. H. and Zubarev, R. A.** (2001). Advantages of external accumulation for electron capture dissociation in Fourier transform mass spectrometry. *Analytical Chemistry* **73**, 2998–3005.
- Haslam, S. M., Coles, G. C., Morris, H. R. and Dell, A.** (2000). Structural characterization of the N-glycans of *Dictyocaulus viviparus*: discovery of the Lewis X structure in a nematode. *Glycobiology* **10**, 223–229.
- Haslam, S. M., Coles, G. C., Munn, E. A., Smith, T. S., Smith, H. F., Morris, H. R. and Dell, A.** (1996). *Haemonchus contortus* glycoproteins contain N-linked oligosaccharides with novel highly fucosylated core structures. *Journal of Biological Chemistry* **271**, 30561–30570.
- Haslam, S. M., Houston, K. M., Harnett, W., Reason, A. J., Morris, H. R. and Dell, A.** (1999). Structural studies of N-glycans of filarial parasites. Conservation of phosphorylcholine-substituted glycans among species and discovery of novel chito-oligomers. *Journal of Biological Chemistry* **274**, 20953–20960.
- Hayes, K. S., Bancroft, A. J. and Grecnis, R. K.** (2004). Immune-mediated regulation of chronic intestinal nematode infection. *Immunological Reviews* **201**, 75–88.
- Hayunga, E. G., Murrell, K. D., Taylor, D. W. and Vannier, W. E.** (1979). Isolation and characterization of surface antigens from *Schistosoma mansoni*. Evaluation of techniques for radioisotope labeling of surface proteins from adult worms. *Journal of Parasitology* **65**, 488–496.
- Hernandez, J. D. and Baum, L. G.** (2002). Ah sweet mystery of death! Galectins and control of cell fate. *Glycobiology* **12**, 127–136.
- Hewitson, J. P., Harcus, Y. M., Curwen, R. S., Dowle, A. A., Atmadja, A. K., Ashton, P. D., Wilson, A. and Maizels, R. M.** (2008). The secretome of the filarial parasite, *Brugia malayi*: proteomic profile of adult excretory-secretory products. *Molecular and Biochemical Parasitology* **160**, 8–21.
- Hill, D. E., Gamble, R., Rhoads, M. L., Fetterer, R. H. and Urban, J. F., Jr.** (1993). *Trichuris suis*: a zinc metalloprotease from culture fluids of adult parasites. *Experimental Parasitology* **77**, 170–178.
- Holland, M. J., Harcus, Y., Balic, A. and Maizels, R. M.** (2005). TH2 induction by *Nippostrongylus* secreted antigens in mice deficient in B cells, eosinophils or MHC Class I-related receptors. *Immunology Letters* **96**, 93–101.
- Holland, M. J., Harcus, Y. M., Riches, P. L. and Maizels, R. M.** (2000). Proteins secreted by the parasitic nematode *Nippostrongylus brasiliensis* act as adjuvants for TH2 responses. *European Journal of Immunology* **30**, 1977–1987.
- Horii, Y., Owhashi, M., Ishii, A., Bandou, K. and Usui, M.** (1984). Eosinophil and neutrophil chemotactic activities of adult worm extracts of *Schistosoma japonicum* *in vivo* and *in vitro*. *Journal of Parasitology* **70**, 955–961.
- Houston, K. M. and Harnett, W.** (2004). Structure and synthesis of nematode phosphorylcholine-containing glycoconjugates. *Parasitology* **129**, 655–661.
- Houston, K. M., Sutharsan, R., Steiger, C. N., Scachter, H. and Harnett, W.** (2008). Gene inactivation confirms the identity of enzymes involved in nematode phosphorylcholine-N-glycan synthesis. *Molecular and Biochemical Parasitology* **157**, 88–91.
- Hsieh, G. C. F., Loukas, A., Wahl, A. M., Bhatia, M., Wang, Y., Williamson, A. L., Kehn, K. W., Maruyama, H., Hotez, P. J., Leitenberg, D., Bethony, J. and Constant, S. L.** (2004). A secreted protein from the human hookworm *Necator americanus* binds selectively to NK cells and induces IFN γ production. *Journal of Immunology* **173**, 2699–2704.
- Huang, Y., Konse, T., Mechref, Y. and Novotny, M. V.** (2002). Matrix assisted desorption ionization mass spectrometry compatible with beta-elimination of O-linked oligosaccharides. *Rapid Communications in Mass Spectrometry* **16**, 1199–1204.
- Huberty, M. C., Vath, J. E., Yu, W. and Martin, S. A.** (1993). Site specific carbohydrate identification in recombinant proteins using MALDI-TOF-MS. *Analytical Chemistry* **65**, 2791–2800.
- Humphreys, N. E., Xu, D., Hepworth, M. R., Liew, F. Y. and Grecnis, R. K.** (2008). IL-33, a potent

- inducer of adaptive immunity to intestinal nematodes. *Journal of Immunology* **180**, 2443–2449.
- Hunter, M. M., Wang, A. and McKay, D. M.** (2007). Helminth infection enhances disease in a murine TH2 model of colitis. *Gastroenterology* **132**, 1320–1330.
- Ikehara, Y., Ikehara, S. K. and Paulson, J. C.** (2004). Negative regulation of T cell receptor signaling by Siglec-7 and Siglec-9. *Journal of Biological Chemistry* **279**, 43117–43125.
- Irwin, J. A., Morrissey, P. E. W., Ryan, J. P., Walshe, A., O'Neill, S. M., Carrington, S. D., Matthews, E., Fitzpatrick, E., Mulcahy, G., Corfield, A. P. and Dalton, J. P.** (2004). Glycosidase activity in the excretory-secretory products of the liver fluke, *Fasciola hepatica*. *Parasitology* **129**, 465–472.
- Itami, D. M., Oshiro, T. M., Araujo, C. A., Perini, A., Martins, M. A., Macedo, M. S. and Macedo-Soares, M. F.** (2005). Modulation of murine experimental asthma by *Ascaris suum* components. *Clinical and Experimental Allergy* **35**, 873–879.
- Jackson, S. K., Abate, W., Parton, J., Jones, S. and Harwood, J. L.** (2008). Lysophospholipid metabolism facilitates Toll-like receptor 4 membrane translocation to regulate the inflammatory response. *Journal of Leukocyte Biology* **84**, 86–92.
- Jefferies, J. R., Turner, R. J. and Barrett, J.** (1997). Effect of *Fasciola hepatica* excretory-secretory products on the metabolic burst of sheep and human neutrophils. *International Journal for Parasitology* **27**, 1025–1029.
- Jenkins, S. J., Hewitson, J. P., Ferret-Bernard, S. and Mountford, A. P.** (2005). Schistosome larvae stimulate macrophage cytokine production through TLR4-dependent and independent pathways. *International Immunology* **17**, 1409–1418.
- Joesph, G. T., Huima, T., Lion, A. and Lustigman, S.** (2000). A novel developmentally regulated galectin of *Onchocerca volvulus*. *Molecular and Biochemical Parasitology* **106**, 187–195.
- Joshi, A. D., Raymond, T., Coelho, A. L., Kunkel, S. L. and Hogaboam, C. M.** (2008). A systemic granulomatous response to *Schistosoma mansoni* eggs alters responsiveness of bone marrow derived macrophages to Toll-like receptor agonists. *Journal of Leukocyte Biology* **83**, 314–324.
- Judson, D. G., Dixon, J. B. and Skerriit, G. C.** (1987). Occurrence and biochemical characteristics of cestode lymphocyte mitogens. *Parasitology* **94**, 151–160.
- Junking, M., Wongkham, C., Sripa, B., Sawanyawisuth, K., Araki, N. and Wongkham, S.** (2008). Decreased expression of galectin-3 is associated with metastatic potential of liver fluke-associated cholangiocarcinoma. *European Journal of Cancer* **44**, 619–626.
- Kang, S., Cummings, R. D. and McCall, J. W.** (1993). Characterization of the N-linked oligosaccharides in glycoproteins synthesized by microfilariae of *Dirofilaria immitis*. *Journal of Parasitology* **79**, 815–828.
- Kanse, S. M., Liang, O., Schubert, U., Haas, H., Preissner, K. T., Doenhoff, M. J. and Dennis, R. D.** (2005). Characterization and partial purification of *Schistosoma mansoni* egg-derived pro-angiogenic factor. *Molecular and Biochemical Parasitology* **144**, 76–85.
- Kasper, G., Brown, A., Eberl, M., Vallar, L., Kieffer, N., Berry, C., Girdwood, K., Eggleton, P., Quinnell, R. and Pritchard, D. I.** (2001). A calreticulin-like molecule from the human hookworm *Necator americanus* interacts with C1q and the cytoplasmic signaling domain of some integrins. *Parasite Immunology* **23**, 141–152.
- Kato, Y. and Komatsu, S.** (1996). ASABF, a novel cysteine rich antibacterial peptide isolated from the nematode *Ascaris suum*: purification, primary structure and molecular cloning of cDNA. *Journal of Biological Chemistry* **271**, 30493–30498.
- Katz, E., Deehan, M. R., Seatter, S., Lord, C., Sturrock, R. D. and Harnett, M. M.** (2001). B cell receptor stimulated mitochondrial phosphorylipase A2 activation and resultant disruption of mitochondrial membrane potential correlate with the induction of apoptosis in WEH 1–231. B cells. *Journal of Immunology* **166**, 137–147.
- Kean, D. E., Ohtsuka, I., Sato, K., Hada, N., Takeda, T., Lochnit, G., Geyer, R., Harnett, M. M. and Harnett, W.** (2006). Dissecting *Ascaris* glycosphingolipids for immunomodulatory moieties – the use of synthetic structural glycosphingolipid analogues. *Parasite Immunology* **28**, 69–76.
- Keir, P. A., Brown, D. M., Clouter-Baker, A., Harcus, Y. M. and Proudfoot, L.** (2004). Inhibition of neutrophil recruitment by ES of *Nippostrongylus brasiliensis*. *Parasite Immunology* **26**, 137–139.
- Kelm, S., Pelz, A., Schauer, R., Filbin, M. T., Tang, S., DeBellard, M. E., Schnaar, R. I., Mahoney, J. A., Hartnell, A., Bradfield, P. and Crocker, P. R.** (1994). Sialoadhesion, MAG and CD22 define a new family of sialic acid dependent adhesion molecules of the immunoglobulin superfamily. *Current Biology* **4**, 965–972.
- Khan, W. I.** (2008). Physiological changes in the gastrointestinal tract and host protective immunity: learning from the mouse-*Trichinella spiralis* model. *Parasitology* **135**, 671–682.
- Khoo, K. H., Sarda, S., Xu, X., Caulfield, J. P., McNeil, M. R., Homans, S. W., Morris, H. R. and Dell, A.** (1995). A unique multifucosylated 3GalNAc β 1,4GlcNAc β 1,3Gal α 1 motif constitutes the repeating unit of the complex O-glycans derived from the cercarial glycocalyx of *Schistosoma mansoni*. *Journal of Biological Chemistry* **270**, 17114–17123.
- Kim, Y. J., Choi, M. H., Hong, S. T. and Bae, Y. M.** (2008). Proliferative effects of excretory-secretory products from *Clonorchis sinensis* on the human epithelial cell line HEK293 via regulation of the transcription factor E2F1. *Parasitology Research* **102**, 411–417.
- Kina, Y., Fukumoto, S., Miura, K., Tadamoto, S., Nunomura, K., Diraghayu, P. and Hirai, K.** (2005). A glycoprotein from *Spirometra erinaceieuropaei* plerocercoids suppresses osteoclastogenesis and proinflammatory cytokine gene expression. *International Journal for Parasitology* **35**, 1399–1406.
- King, C. H. and Dangerfield-Cha, M.** (2008). The unacknowledged impact of chronic schistosomiasis. *Chronic Illness* **4**, 65–79.
- Klion, A. D. and Donelson, J. E.** (1994). OvGalBP, a filarial antigen with homology to vertebrate

- galactoside-binding proteins. *Molecular and Biochemical Parasitology* **65**, 305–315.
- Ko, A. I., Drager, U. C. and Harn, D. A.** (1990). A *Schistosoma mansoni* epitope recognized by a protective monoclonal antibody is identical to the stage specific embryonic antigen 1 (SSEA-1). *Proceedings of the National Academy of Sciences, USA* **87**, 4159–4163.
- Kooyman, F. N. J., de Vries, E., Ploegar, H. W. and van Putten, J. P. M.** (2007). Antibodies elicited by the bovine lungworm, *Dictyocaulus viviparus*, cross-react with platelet activating factor. *Infection and Immunity* **75**, 4456–4462.
- Kottgen, M., Loffler, T., Jacobi, C., Nitchke, R., Pavenstadt, H., Schreiber, R., Frische, S., Nielsen, S. and Leipziger, J.** (2003). P2Y₆ receptor mediates colonic NaCl secretion via differential activation of cAMP-mediated transport. *Journal of Clinical Investigation* **111**, 371–379.
- Kradin, R. L., Badizadegan, K., Auluck, P., Korzenik, J. and Lauwers, G. Y.** (2006). Iatrogenic *Trichuris suis* infection in a patient with Crohn's disease. *Archives of Pathology and Laboratory Medicine* **130**, 718–720.
- Kron, M., Marquard, K., Hartlein, M., Price, S. and Leberman, R.** (1995). An immunodominant antigen of *Brugia malayi* is an asparaginyl-tRNA synthetase. *FEBS Letters* **374**, 122–124.
- Kumar, S. and Pritchard, D. I.** (1992). The partial characterization of proteases present in the excretory-secretory products and exsheathing fluid of the infective L3 larva of *Necator americanus*. *International Journal for Parasitology* **22**, 563–572.
- Lajaunias, F., Dayer, J. M. and Chizzolini, C.** (2005). Constitutive repressor activity of CD33 on human monocytes requires sialic acid recognition and phosphoinositide 3 kinase-mediated intracellular signaling. *European Journal of Immunology* **35**, 243–251.
- Lee, J. C., Cho, G. S., Kwon, J. H., Shin, M. H., Lim, J. H. and Kim, W. K.** (2006). Macrophageal/microglial cell activation and cerebral injury induced by excretory-secretory products secreted by *Paragonimus westermani*. *Neuroscience Research* **54**, 133–139.
- Leffler, H. and Barondes, S. H.** (1986). Specificity of binding of three soluble rat lung lectins to substituted and unsubstituted mammalian beta-galactosides. *Journal of Biological Chemistry* **261**, 10119–10126.
- Leid, R., Suquet, C. M. and Perryman, L. E.** (1984). Inhibition of antigen and lectin induced proliferation of rat spleen cells by *Taenia taeniformes* proteinase inhibitor. *Clinical and Experimental Immunology* **57**, 187–194.
- Leid, R. W., Suquet, C. M., Bouwer, H. G. A. and Hinrichs, D. J.** (1986). Interleukin inhibition by a parasite proteinase inhibitor, taeniaestatin. *Journal of Immunology* **137**, 2700–2702.
- Liaudet, L., Mabley, J. G., Pacher, P., Virag, L., Soriano, F. G., Marton, A., Hasko, G., Deitch, E. A. and Szabo, C.** (2002). Inosine exerts a broad range of anti-inflammatory effects in a murine model of acute lung injury. *Annals in Surgery* **235**, 568–578.
- Lima, C., Perini, A., Garcia, M. L. B., Martins, M. A., Teixeira, M. M. and Macedo, M. S.** (2002). Eosinophilic inflammation and airway hyper-responsiveness are profoundly inhibited by a helminth (*Ascaris suum*) extract in a murine model of asthma. *Clinical and Experimental Allergy* **32**, 1659–1666.
- Loeffler, D. A., Lunfy, S. K., Singh, K. P., Gerard, H. C., Hudson, A. P. and Boros, D. L.** (2002). Soluble egg antigens from *Schistosoma mansoni* induce angiogenesis-related processes by up-regulating vascular endothelial growth factor in human endothelial cells. *Journal of Infectious Disease* **185**, 1650–1656.
- Loukas, A., Doedens, A., Hintz, M. and Maizels, R. M.** (2000a). Identification of a new C-type lectin, TES-70, secreted by infective larvae of *Toxocara canis*, which binds to host ligands. *Parasitology* **121**, 545–554.
- Loukas, A., Hintz, M., Linder, D., Mullin, N. P., Parkinson, J., Tetteh, K. K. A. and Maizels, R. M.** (2000b). A family of secreted mucins from the parasitic nematode *Toxocara canis* bears diverse mucin domains but shares similar flanking six cysteine repeat motifs. *Journal of Biological Chemistry* **275**, 39600–39607.
- Loukas, A., Mullin, N. P., Tetteh, K. K. A., Moens, L. and Maizels, R. M.** (1999). A novel C-type lectin secreted by a tissue dwelling parasitic nematode. *Current Biology* **9**, 825–828.
- Lundy, S. K., Lerman, S. P. and Boros, D. L.** (2001). Soluble egg antigen stimulated T Helper lymphocyte apoptosis and evidence for cell death mediated by FasL⁺ T and B cells during murine *Schistosoma mansoni* infection. *Infection and Immunity* **69**, 271–280.
- Mabley, J. G., Pacher, P., Liaudet, L., Soriano, F. G., Hasko, G., Marton, A., Szabo, C. and Salzman, A. L.** (2003). Inosine reduces inflammation and improves survival in a murine model of colitis. *American Journal of Physiology Gastrointestinal and Liver Physiology* **284**, 138–144.
- Maizels, R. M., Balic, A., Gomez-Escobar, N., Nair, M., Taylor, M. D. and Allen, J. E.** (2004). Helminth parasites – masters of regulation. *Immunology Reviews* **201**, 89–116.
- Marshall, A. J., Brunet, L. R., van Dessel, Y., Alcaez, A., Bliss, S. K., Pearce, E. J. and Denkers, E. Y.** (1999). *Toxoplasma gondii* and *Schistosoma mansoni* synergize to promote hepatocyte dysfunction associated with high levels of plasma TNF α and early death in C57BL/6 mice. *Journal of Immunology* **163**, 2089–2097.
- Marsland, B. J., Camberis, M. and Le Gros, G.** (2005). Secretory products from infective forms of *Nippostrongylus brasiliensis* induce a rapid allergic airway inflammatory response. *Immunology and Cell Biology* **83**, 40–47.
- McConchie, B. W., Norris, H. H., Bundoc, V. G., Trivedi, S., Boesen, A., Urban, J. F., Jr. and Keane-Meyers, A. M.** (2006). *Ascaris suum*-derived products suppress mucosal allergic inflammation in an interleukin 10 independent manner via interference with dendritic cell function. *Infection and Immunity* **74**, 6632–6641.
- McDiarmid, S. S. and Podesta, R. B.** (1984). Identification of a sialic acid containing glycocalyx on the surface of *Schistosoma mansoni*. *Molecular and Biochemical Parasitology* **10**, 33–43.
- McInnes, I. B., Leung, B. P., Harnett, M., Gracie, J. A., Liew, F. Y. and Harnett, W.** (2003). A novel therapeutic approach targeting articular inflammation using the filarial nematode-derived phosphorylcholine-containing glycoprotein ES-62. *Journal of Immunology* **171**, 2127–2133.

- McKay, D. M.** (2006). The beneficial helminth parasite? *Parasitology* **132**, 1–12.
- McKerrow, J. H., Brindley, P., Brown, M., Gam, A. A., Staunton, C. and Neva, F. A.** (1990). *Strongyloides stercoralis*: identification of a protease that facilitates penetration of skin by the infective larvae. *Experimental Parasitology* **70**, 134–143.
- Melendez, A. J., Harnett, M. M., Pushparaj, P. N., Wong, W. S., Tay, H. K., McSharry, C. P. and Harnett, W.** (2007). Inhibition of Fc epsilon RI-mediated mast cell responses by ES-62, a product of parasitic filarial nematodes. *Nature Medicine* **13**, 1375–1381.
- Meyer, S., van Liempt, E., Imberty, A., van Kooyk, Y., Geyer, H., Geyer, R. and van Die, I.** (2005). DC-SIGN mediates binding of dendritic cells to authentic pseudo-LewisY glycolipids of *Schistosoma mansoni* cercariae, the first parasite specific ligand of DC-SIGN. *Journal of Biological Chemistry* **280**, 37349–37359.
- Mikes, L. and Horak, P.** (2001). A protein with lectin activity in the penetration glands of *Diplostomum pseudopathaceum* cercariae, *International Journal for Parasitology* **31**, 245–252.
- Milos, N. and Zalik, S. E.** (1983). Calcium-independent adhesion of extra-embryonic endoderm cells from the early chick blastoderm is inhibited by the blastoderm beta-D-galactoside-binding lectin and by beta-galactosidase. *Cell Differentiation* **12**, 341–347.
- Morelle, W. and Michaelski, J. C.** (2004). The mass spectrometric analysis of glycoproteins and their glycan structures. *Current Analytical Chemistry* **1**, 29–57.
- Morris, H. R., Paxton, T., Panico, M., McDowell, R. and Dell, A.** (1997). A novel geometry mass spectrometer, the Q-TOF, for low femtomole/attomole range biopolymer sequencing. *Journal of Protein Chemistry* **16**, 469–479.
- Moyle, M., Foster, D. L., McGrath, D. E., Brown, S. M., Laroche, Y., De Meutter, J., Stanssens, P., Bogowitz, C. A., Fried, V. A., Ely, J. A., Soule, H. R. and Vlasuk, G. P.** (1994). A hookworm glycoprotein that inhibits neutrophil function is a ligand of the integrin CD11b/CD18. *Journal of Biological Chemistry* **269**, 10008–10015.
- Nagy, K., Vekey, K., Imre, T., Ludanyi, K., Barrow, M. P. and Derrick, P. J.** (2004). Electrospray ionization fourier transform ion cyclotron resonance mass spectrometry of human alpha-1-acid glycoprotein. *Analytical Chemistry* **76**, 4998–5005.
- Nutku, E., Aizawa, H., Hudson, S. A. and Bochner, B. S.** (2003). Ligation of Siglec-8: selective mechanism for induction of human eosinophil apoptosis. *Blood* **101**, 5014–5020.
- Nyame, K., Cummings, R. D. and Damian, R. T.** (1986). Characterization of the O-linked oligosaccharides in glycoproteins synthesized by *Schistosoma mansoni* schistosomula. *Journal of Parasitology* **74**, 562–572.
- Nyame, K., Smith, D. F., Damian, R. T. and Cummings, R. D.** (1989). Complex type asparagine-linked oligosaccharides in glycoproteins synthesized by *Schistosoma mansoni* adult males contain terminal beta-linked N-acetylgalactosamine. *Journal of Biological Chemistry* **264**, 3235–3243.
- Oberhelman, R. A., Guerrero, E. S., Fernandez, M. L., Silio, M., Mercado, D., Comiskey, N., Ihenacho, G. and Mera, R.** (1998). Correlations between intestinal parasitosis, physical growth and psychomotor development in infants and children from rural Nicaragua. *American Journal of Tropical Medicine* **58**, 470–475.
- O'Neill, S. M., Mills, K. H. and Dalton, J. P.** (2001). *Fasciola hepatica* cathepsin L cysteine proteinase suppresses *Bordetella pertussis*-specific interferon gamma production *in vivo*. *Parasite Immunology* **23**, 541–547.
- Okano, M., Satoskar, A. R., Nishizaki, K., Abe, M. and Harn, D. A.** (1999). Induction of TH2 responses and IgE is largely due to carbohydrates functioning as adjuvants on *Schistosoma mansoni* egg antigens. *Journal of Immunology* **163**, 6712–6717.
- Osborne, J. and Devaney, E.** (1999). Interleukin-10 and antigen presenting cells actively suppress Th1 cells in BALB/c mice infected with the filarial parasite *Brugia pahangi*. *Infection and Immunity* **67**, 1509–1605.
- Owhashi, M. and Ishii, A.** (1982). Purification and characterization of a high molecular weight eosinophil chemotactic factor from *Schistosoma japonicum* eggs. *Journal of Immunology* **129**, 2226–2231.
- Pastrana, D. V., Raghavan, N., Fitzgerald, P., Eisinger, S. W., Metz, C., Bucala, R., Schleimer, R. P., Bickel, C. and Scott, A. L.** (1998). Filarial nematode parasites secrete a homologue of the human cytokine macrophage migration inhibitory factor. *Infection and Immunity* **66**, 5955–5963.
- Perrigou, J. G., Marshall, F. and Artis, D.** (2008). On the hunt for helminths: innate immune cells in the recognition and response to helminth parasites. *Cellular Microbiology* **10**, 1757–1764.
- Persat, F., Vincent, C., Schmitt, D. and Mojon, M.** (1996). Inhibition of human peripheral blood mononuclear cell proliferative response by glycosphingolipids from metacestodes of *Echinococcus multilocularis*. *Infection and Immunity* **64**, 3682–3687.
- Persuad, R., Wang, A., Reardon, C. and McKay, D. M.** (2007). Characterisation of the immunoregulatory response to rat the tapeworm *Hymenolepis diminuta* in the non-permissive mouse host. *International Journal for Parasitology* **37**, 393–403.
- Pillai, A., Ueno, S., Zhang, H., Lee, J. M. and Kato, Y.** (2005). Cecropin P1 and novel cecropins: a bacteria-inducible antimicrobial peptide family in the nematode *Ascaris suum*. *The Biochemical Journal* **390**, 207–214.
- Plows, L. D., Cook, R. T., Davies, A. J. and Walker, A. J.** (2005). Carbohydrates that mimic schistosome surface coat components affect ERK and PKC signaling in *Lymnaea stagnalis* haemocytes. *International Journal for Parasitology* **35**, 293–302.
- Powell, L. D. and Varki, A.** (1994). The oligosaccharide binding specificities of CD22-beta, a sialic acid specific lectin of B cells. *Journal of Biological Chemistry* **269**, 10628–10636.
- Pyz, E., Marshall, A. S. J., Gordon, S. and Brown, G. D.** (2006). C-type lectin-like receptors on myeloid cells. *Annals of Medicine* **38**, 242–251.

- Rabinovich, G. A., Liu, F. T., Hirashima, M. and Anderson, A.** (2007). An emerging role for galectins in tuning the immune response: lessons from experimental models of inflammatory disease, autoimmunity and cancer. *Scandinavian Journal of Immunology* **66**, 143–158.
- Rabinovich, G. A., Baum, L. G., Tinari, N., Paganelli, R., Natoli, C., Liu, F. T. and Iacobelli, S.** (2002). Galectins and their ligands: amplifiers, silencers or tuners of the inflammatory response? *Trends in Immunology* **23**, 313–320.
- Rakha, N. K., Dixon, J. B., Carter, S. D., Craig, P. S., Jenkins, P. and Folkard, S.** (1991). *Echinococcus multilocularis* antigens modify accessory cell function of macrophages. *Immunology* **74**, 652–656.
- Ramajo-Hernandez, A., Olega, A., Ramajo-Martin, V. and Perez-Sanchez, R.** (2007). Carbohydrate profiling and protein identification of tegumental and excreted/secreted glycoproteins of adult *Schistosoma bovis* worms. *Veterinary Parasitology* **144**, 45–60.
- Ramirez, B. L., Zack Howard, O. M., Fang Don, H., Edamatsu, T., Hartlein, M. and Kron, M.** (2006). *Brugia malayi* asparaginyl-transfer RNA synthetase induces chemotaxis of human leukocytes and activates G-protein-coupled receptors CXCR1 and CXCR2. *Journal of Infectious Diseases* **193**, 1164–1171.
- Rathore, D. K., Suchitra, S., Suini, M., Singh, B. P. and Joshi, P.** (2006). Identification of a 66 kDa *Haemonchus contortus* excretory/secretory antigen inhibiting host monocytes. *Veterinary Parasitology* **138**, 291–300.
- Reese, T. A., Liang, H. E., Tager, A. M., Luster, A. D., van Rooijen, N., Voehringer, D. and Locksley, R. M.** (2007). Chitin induces accumulation in tissue of innate immune cells associated with allergy. *Nature Letters* **447**, 92–97.
- Reyes, J. L. and Terrazas, L. I.** (2007). The divergent roles of alternatively activated macrophages in helminthic infections. *Parasite Immunology* **29**, 609–619.
- Rigano, R., Profumo, E., Brushi, F., Carulli, G., Azzara, A., Ioppolo, S., Buttari, B., Ortona, E., Margutti, P., Teggi, A. and Siracusano, A.** (2001). Modulation of human immune response by *Echinococcus granulosus* antigen B and its possible role in evading host defences. *Infection and Immunity* **69**, 288–296.
- Rigano, R., Buttari, B., Profumo, E., Ortona, E., Delunardo, F., Margutti, P., Mattei, V., Teggi, A., Sorice, M. and Siracusano, A.** (2007). *Echinococcus granulosus* antigen B impairs human dendritic cell differentiation and polarizes immature dendritic cell maturation towards a TH2 cell response. *Infection and Immunity* **75**, 1667–1678.
- Rikihisa, Y., Lin, Y. C. and Fukaya, T.** (1985). *Taenia taeniformes*: inhibition of rat testosterone production by excretory-secretory product of the cultured metacystode. *Experimental Parasitology* **59**, 390–397.
- Rocha, F. A. C., Leite, A. K. R. M., Pompeu, M. M. L., Cunha, T. M., Verri, W. A., Jr., Soares, F. M., Castro, R. R. and Cunha, F. Q.** (2008). Protective effect of an extract from *Ascaris suum* in experimental arthritis models. *Infection and Immunity* **76**, 2736–2745.
- Salatino, M., Croci, D. O., Bianco, G. A., Illarregui, G. M., Toscano, M. A. and Rabinovich, G. A.** (2008). Galectin-1 as a potential therapeutic target in autoimmune disorders and cancer. *Expert Opinion on Biological Therapy* **8**, 45–57.
- Schallig, H. D. F. G. and van Leeuwen, M. A. W.** (1996). Carbohydrate epitopes on *Haemonchus contortus* antigens. *Parasitology Research* **82**, 38–42.
- Schnoeller, C., Rausch, S., Pillai, S., Avagyan, A., Wittig, B. M., Loddenkemper, C., Hamann, A., Hamelmann, E., Lucius, R. and Hartmann, S.** (2008). A helminth immunomodulator reduces allergic and inflammatory responses by induction of IL-10-producing macrophages. *Journal of Immunology* **180**, 4265–4272.
- Schonemeyer, A., Lucius, R., Sonnenburg, B., Brattig, N., Sabat, R., Schilling, K., Bradley, J. and Hartmann, S.** (2001). Modulation of human T cell responses and macrophage functions by onchocystatin, a secreted protein of the filarial nematode *Onchocerca volvulus*. *Journal of Immunology* **167**, 3207–3215.
- Scuitto, E., Fragoso, G., Baca, M., de la Cruz, V., Lemus, L. and Lamoyi, E.** (1995). Depressed T-cell proliferation associated with susceptibility to experimental *Taenia crassiceps* infection. *Infection and Immunity* **63**, 2277–2281.
- Segura, M., Su, Z., Piccirillo, C. and Stevenson, M. M.** (2007). Impairment of dendritic cell function by excretory-secretory products: a potential mechanism for nematode-induced immunosuppression. *European Journal of Immunology* **37**, 1887–1904.
- Serradell, M. C., Guasconi, L., Cervi, L., Chiapello, L. S. and Masih, D. T.** (2007). Excretory-secretory products from *Fasciola hepatica* induce eosinophil apoptosis by a caspase-dependent mechanism. *Veterinary Immunology and Immunopathology* **117**, 197–208.
- Shepherd, J. C., Aitken, A. and McManus, D. P.** (1991). A protein secreted *in vivo* by *Echinococcus granulosus* inhibits elastase activity and neutrophil chemotaxis. *Molecular and Biochemical Parasitology* **44**, 81–90.
- Smith, A. M., Dowd, A. J., Heffernan, M., Robertson, C. D. and Dalton, J. P.** (1993). *Fasciola hepatica*: a secreted cathepsin L-like proteinase cleaves host immunoglobulin. *International Journal for Parasitology* **23**, 977–983.
- Souza, V. M. O., Faquin-Mauro, E. L. and Macedo, M. S.** (2002). Extracts of *Ascaris suum* egg and adult worm share similar immunosuppressive properties. *Brazilian Journal of Medical and Biological Research* **35**, 81–89.
- Spolski, R. J., Corson, J., Thomas, P. G. and Kuhn, R. E.** (2000). Parasite secreted products regulate the host response to larval *Taenia crassiceps*. *Parasite Immunology* **22**, 297–305.
- Spolski, R. J., Thomas, P. G., See, E. J., Mooney, K. A. and Kuhn, R. E.** (2002). Larval *Taenia crassiceps* secretes a protein with characteristics of murine interferon- γ . *Parasitology Research* **88**, 431–438.
- Srivatsan, J., Smith, D. F. and Cummings, R. D.** (1992). The human blood fluke *Schistosoma mansoni* synthesizes glycoproteins containing the LewisX antigen. *Journal of Biological Chemistry* **267**, 20196–20203.

- Stein, M., Keshav, S., Harris, N. and Gordon, S.** (1992). Interleukin 4 potently enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. *Journal of Experimental Medicine* **176**, 287–292.
- Stepek, G., Auchie, M., Tate, R., Watson, K., Russell, D. G., Devaney, E. and Harnett, W.** (2002). Expression of the filarial nematode phosphorylcholine-containing glycoprotein, ES-62, is stage specific. *Parasitology Today* **125**, 155–164.
- Stepek, G., Houston, K. M., Goodridge, H. S., Devaney, E. and Harnett, W.** (2004). Stage specific and species specific differences in the production of the mRNA and protein for the filarial nematode secreted product ES-62. *Parasitology Today* **128**, 91–98.
- Suchitra, S. and Joshi, P.** (2005). Characterization of *Haemonchus contortus* calreticulin suggests its role in feeding and immune evasion by the parasite. *Biochimica et Biophysica Acta* **1722**, 293–303.
- Suchitra, S., Anbu, K. A., Rathore, D. K., Mahawar, M., Singh, B. P. and Joshi, P.** (2008). *Haemonchus contortus* calreticulin binds to C-reactive protein of its host, a novel survival strategy of the parasite. *Parasite Immunology* **30**, 371–374.
- Summers, R. W., Elliott, D. E., Qadir, K., Urban, J. F., Jr., Thompson, R. and Weinstock J. V.** (2003). *Trichuris suis* seems to be safe and possibly effective in the treatment of inflammatory bowel disease. *American Journal of Gastroenterology* **98**, 2034–2041.
- Summers, R. W., Elliott, D. E., Urban, J. F., Jr., Thompson, R. A. and Weinstock, J. V.** (2005). *Trichuris suis* therapy for active ulcerative colitis: a randomized control trial. *Gastroenterology* **128**, 825–832.
- Suquet, C. M., Green-Edwards, C. and Leid, R. W.** (1984). Isolation and partial characterization of a *Taenia taeniformes* metacystode proteinase inhibitor. *International Journal for Parasitology* **14**, 165–172.
- Suttiprapa, S., Loukas, A., Laha, T., Wongkham, S., Kaewkes, S., Gaze, S., Brindley, P. J. and Sripa, B.** (2008). Characterization of the anti-oxidant enzyme, thioredoxin peroxidase, from the carcinogenic human liver fluke, *Opisthorchis viverrini*. *Molecular and Biochemical Parasitology* **160**, 116–122.
- Tanaka, J. and Torisu, M.** (1978). *Anisakis* and eosinophil. Detection of a soluble factor selectively chemotactic for eosinophils in the extract from *Anisakis* larvae. *Journal of Immunology* **120**, 745–749.
- Tanaka, J., Baba, T. and Torisu, M.** (1979). *Ascaris* and eosinophil. Isolation and characterization of eosinophil chemotactic factor and neutrophil chemotactic factor of parasite in *Ascaris* antigen. *Journal of Immunology* **122**, 302–308.
- Taylor, K. and Hoole, D.** (1997). Interactions between rat C-reactive protein and adult *Hymenolepis diminuta*. *Parasitology* **115**, 297–302.
- Taylor, J. J., Mohrs, M. and Pearce, E. J.** (2006). Regulatory T cells responses develop in parallel to Th responses and control the magnitude and phenotype of the Th effector population. *Journal of Immunology* **176**, 5839–5847.
- Terrazas, L. I., Walsh, K. L., Piskorska, D., McGuire, E. and Harn, D. A.** (2001). The schistosome oligosaccharide lacto-N-neotetrose expands Gr1⁺ cells that secrete anti-inflammatory cytokines and inhibit proliferation of naive CD4⁺ cells: a potential mechanism for immune polarization in helminth infections. *Journal of Immunology* **167**, 5294–5303.
- Thomas, P. G., Carter, M. R., Atochina, O., Da'Dara, A. A., Piskorska, D., McGuire, E. and Harn, D. A.** (2003). Maturation of dendritic cell 2 phenotype by a helminth glycan uses a Toll-like receptor 4 dependent mechanism. *Journal of Immunology* **171**, 5837–5841.
- Thomas, P. G., Carter, M. R., Da'Dara, A. A., DeSimone, T. M. and Harn, D. A.** (2005). A helminth glycan induces APC maturation via alternative NFκB activation independent of IκBα degradation. *Journal of Immunology* **175**, 2082–2090.
- Trujillo-Vargas, C. M., Werner-Klein, M., Wohlleben, G., Polte, T., Hansen, G., Ehlers, S. and Erb, K. J.** (2007). Helminth-derived products inhibit the development of allergic responses in mice. *American Journal of Respiratory and Critical Care Medicine* **175**, 336–344.
- Truong, M. J., Gruart, V., Liu, F. T., Prin, L., Capron, A. and Capron, M.** (1993). IgE-binding molecules (Mac-2εBP) expressed by human eosinophils: implication in IgE-dependent eosinophil cytotoxicity. *European Journal of Immunology* **23**, 3230–3235.
- Turner, D. G., Wildblood, L. A., Inglis, N. F. and Jones, D. G.** (2008). Characterization of galectin like activity from the parasitic nematode, *Haemonchus contortus*, which modulates ovine eosinophil migration *in vitro*. *Veterinary Immunology and Immunopathology* **122**, 138–145.
- van de Vijver, K. K., Deelder, A. M., Jacobs, W., van Marck, E. A. and Hokke, C. H.** (2006). LacdiNAc- and LacNAc-containing glycans induce granulomas in an *in vivo* model for schistosome egg-induced hepatic granuloma formation. *Glycobiology* **16**, 237–243.
- van den Berg, T. K., Honing, H., Franke, N., van Remoortere, A., Shiphorst, W. E. C. M., Liu, F. T., Deedler, A. M., Cummings, R. D., Hokke, C. H. and van Die, I.** (2004). LacdiNAc-glycans constitute a parasite pattern for galectin-3 mediated immune recognition. *Journal of Immunology* **173**, 1902–1907.
- van der Kleij, D., Latz, E., Brouwers, J. F. H. M., Kruize, Y. C. M., Schmitz, M., Kurt-Jones, E. A., Espevik, T., de Jong, E. C., Kaspenberg, M. L., Golenbock, D. T., Tielens, A. G. M. and Yazdanbakhsh, M.** (2002). A novel host-parasite lipid cross talk: Schistosomal lyso-phosphatidylserine activates Toll-like receptor 2 and affects immune polarization. *Journal of Biological Chemistry* **277**, 48122–48129.
- van Liempt, E., van Vliet, S. J., Engering, A., Vallejo, J. J. G., Bank, C. M. C., Sanchez-Hernandez, M., van Kooyk, Y. and van Die, I.** (2007). *Schistosoma mansoni* soluble egg antigens are internalized by human dendritic cells through multiple C-type lectins and suppress TLR-induced dendritic cell activation. *Molecular Immunology* **44**, 2605–2615.
- van Vliet, S. J., van Liempt, E., Saeland, E., Aarnoudse, C. A., Applemelk, B., Irimura, T., Geijtenbeek, T. B. H., Blijst, O., Alvarez, R., van Die, I. and van Kooyk, Y.** (2005). Carbohydrate profiling reveals a distinctive role for the C-type lectin MGL in the recognition of helminth parasites and

- tumor antigens by dendritic cells. *International Immunology* **17**, 661–669.
- Velupillai, P. and Harn, D. A.** (1994). Oligosaccharide specific induction of interleukin 10 production by B220⁺ cells from schistosome infected mice: a mechanism for regulation of CD4⁺ T cell subsets. *Proceedings of the National Academy of Sciences, USA* **91**, 18–22.
- Velupillai, P., dos Reis, E. A., dos Reis, M. G. and Harn, D. A.** (2000). Lewis X containing oligosaccharide attenuates schistosome egg antigen induced immune depression in human schistosomiasis. *Human Immunology* **61**, 225–232.
- von Gunten, S., Yousefi, S., Seitz, M., Jakob, S. M., Schaffner, T., Seger, R., Takala, J., Villiger, P. M. and Simon, H. U.** (2005). Siglec-9 transduces apoptosis and nonapoptotic death signals into neutrophils depending on the proinflammatory cytokine environment. *Blood* **106**, 1423–1431.
- Wang, A. and McKay, D. M.** (2005). Immune modulation by a high molecular weight fraction from the rat tapeworm, *Hymenolepis diminuta*. *Parasitology* **130**, 575–585.
- Wardlaw, A. C., Forsyth, L. M. G. and Crompton, D. W. T.** (1994). Bactericidal activity in the pig roundworm *Ascaris suum*. *Journal of Applied Bacteriology* **76**, 36–41.
- Weinstock, J. V., Summers, R. W. and Elliott, D. E.** (2005). Role of helminths in regulating mucosal inflammation. *Springer Seminar on Immunology* **27**, 249–271.
- Whelan, M., Harnett, M. M., Houston, K. M., Patel, V., Harnett, W. and Rigley, K. P.** (2000). A filarial nematode secreted product signals dendritic cells to acquire a phenotype that drives the development of TH2 cells. *Journal of Immunology* **164**, 6453–6460.
- Wilson, E. H., Deehan, M. R., Katz, E., Brown, K. S., Houston, K. M., O'Grady, J., Harnett, M. M. and Harnett, W.** (2003). Hyporesponsiveness of murine B lymphocytes exposed to the filarial nematode secreted product ES-62 *in vivo*. *Immunology* **109**, 238–245.
- Wisniewski, N., McNeil, M., Greive, R. B. and Wassom, D. L.** (1993). Characterization of novel fucosylated and tyvelosyl-containing glycoconjugates from *Trichinella spiralis* muscle stage larvae. *Molecular and Biochemical Parasitology* **61**, 25–35.
- Wu, Z., Boonmars, T., Nagano, I., Nakada, T. and Takahashi, Y.** (2003). Molecular expression and characterization of a homologue of host cytokine macrophage migration inhibitory factor from *Trichinella* spp. *Journal of Parasitology* **89**, 507–515.
- Wuhrer, M., Balog, C. I. A., Catalina, M. I., Jones, F. M., Schramm, G., Haas, H., Doenhoff, M. J., Dunne, D. W., Deelder, A. M. and Hokke, C. H.** (2006). IPSE/alpha-1, a major secretory glycoprotein antigen from schistosome eggs, expresses the Lewis X motif on core difucosylated N-glycans. *FEBS Letters* **273**, 2276–2293.
- Wuhrer, M., Kantelhardt, S. R., Dennis, R. D., Doenhoff, M. J., Lochnit, G. and Geyer, R.** (2002). Characterization of glycosphingolipids from *Schistosoma mansoni* eggs carrying Fuc(α1-3)GalNac, GalNac(β1-4)[Fuc(α1-3)]GlcNAc and Gal(β1-4)[Fuc(α1-3)]GlcNAc (Lewis X) terminal structures. *European Journal of Biochemistry* **269**, 481–493.
- Wuhrer, M., Grimm, C., Dennis, R. D., Idris, M. A. and Geyer, R.** (2004). The parasitic trematode *Fasciola hepatica* exhibits mammalian type glycolipids as well as Gal(β1,6)Gal-terminating glycolipids that account for cestode serological cross reactivity. *Glycobiology* **14**, 115–126.
- Yang, R. Y., Hsu, D. K. and Liu, F. T.** (1996). Expression of galectin-3 modulates T cell growth and apoptosis. *Proceedings of the National Academy of Sciences, USA* **93**, 6737–6742.
- Zang, X., Yazdanbakhsh, M., Jiang, H., Kanost, M. R. and Maizels, R. M.** (1999). A novel serpin expressed by blood borne microfilariae of the parasitic nematode *Brugia malayi* inhibits human neutrophil serine proteases. *Blood* **94**, 1418–1428.
- Zhang, J. Q., Nicoll, G., Jones, C. and Crocker, P. R.** (2000). Siglec-9, a novel sialic acid binding member of the immunoglobulin superfamily expressed broadly on human blood leukocytes. *Journal of Biological Chemistry* **275**, 22121–22126.